

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Equidistribution for sets which are not necessarily Galois stable: On a theorem of Mignotte

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1948423> since 2025-01-10T19:43:49Z

Published version:

DOI:10.2422/2036-2145.202309_021

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Equidistribution for sets which are not necessarily Galois stable: On a theorem of Mignotte

Francesco AMOROSO⁽¹⁾ and Arnaud PLESSIS⁽²⁾

⁽¹⁾*Dipartimento di Matematica, Università di Torino, Via Carlo Alberto 10, 10123 Torino, Italia (francesco.amoroso@unito.it)*

⁽²⁾*Yanqi Lake Beijing Institute of Mathematical Sciences and Applications, Huairou District, Beijing 101408, China (plessis@bimsa.cn)*

Abstract

An important result of Bilu deals with the equidistribution of the Galois orbits of a sequence $(\alpha_n)_n$ in $\overline{\mathbb{Q}}^*$. Here, we prove a quantitative equidistribution theorem for a sequence of finite subsets in $\overline{\mathbb{Q}}^*$ which are not necessarily stable by Galois action. We follow a method of Mignotte.

AMS Classification: 11J68, 11G50

1 Introduction

Let X be a metric space. For a finite subset $T \subset X$, the discrete probability measure on X associated to it is given by

$$\mu_{T,X} = \frac{1}{|T|} \sum_{\alpha \in T} \delta_{\alpha,X},$$

where $|T|$ denotes the cardinality of T and $\delta_{\alpha,X}$ the Dirac measure on X supported at α . In the special case $X = \mathbb{C}^*$, we put $\mu_T = \mu_{T,\mathbb{C}^*}$.

We say that a sequence $(\mu_n)_n$ of probability measures on X *converges in distribution* to μ if for every bounded continuous function $f : X \rightarrow \mathbb{C}$, we have

$$\lim_{n \rightarrow +\infty} \int_X f d\mu_n = \int_X f d\mu.$$

An important example of such a sequence was given by Bilu, see Theorem 1.1 below. If the limit above holds for all compactly supported continuous functions, we say that μ_n *weakly converges* to μ .

Throughout this text, we define $h : \overline{\mathbb{Q}} \rightarrow \mathbb{R}$ to be the (logarithmic, absolute) Weil height and λ to be the Haar probability measure on the complex unit circle. We also denote by μ_∞ the set of roots of unity in $\overline{\mathbb{Q}}$.

Theorem 1.1 (Bilu, [9]). *Let K be a number field, and let $(\alpha_n)_n$ be a sequence of $\overline{\mathbb{Q}}^*$ such that $h(\alpha_n) \rightarrow 0$ and $[K(\alpha_n) : K] \rightarrow +\infty$. Then μ_{S_n} converges in distribution to λ , where S_n is the Galois orbit of α_n over K .*

This theorem (which was originally formulated with $K = \mathbb{Q}$) was inspired by a previous work of Szpiro, Ullmo and Zhang who studied the equidistribution of points of small Néron-Tate height on abelian varieties [27]. These two well-known results have been largely generalized to other heights and places, see for instance [30, 25, 5, 7, 23, 17, 8, 12, 28, 19, 13, 11, 14, 6]. Roughly speaking, each one of these results contains an equidistribution theorem *à la* Bilu, that is, a statement of the form “Let K be a number field, and let S_n be a $\text{Gal}(\overline{K}/K)$ -invariant subset of X such that the average of height of α , with $\alpha \in S_n$, goes to 0 and $|S_n| \rightarrow +\infty$. Then $\mu_{S_n, X}$ weakly converges to some Haar measure”.

Here we prove an equidistribution theorem for a sequence of finite subsets in $\overline{\mathbb{Q}}^*$ which are not necessarily stable by Galois action. After posting this paper on ArXiv, Fili informed us that the qualitative version (but not the quantitative one) of our result partially follows from a recent preprint [15], of which we were unaware.

The first avatar of equidistribution theorems is a result of Langevin [20]: Given an open set $\Delta \subset \mathbb{C}$ intersecting the unit circle, the algebraic integers whose Galois conjugates all lie outside Δ cannot have a Weil height arbitrarily small. The idea of Langevin’s proof is to show that the transfinite diameter of the complement of Δ in the unit disk has transfinite diameter less than 1. He then concludes by applying a theorem of Fekete [18] which asserts that there are only a finite number of algebraic integers whose Galois conjugates all belong to a set of transfinite diameter < 1 .

Soon after, Mignotte [22] gives an entirely different proof of Langevin’s result, see [21, Chapter 15] for an excellent expository text. Equidistribution theorems, stated in the modern language of weak convergence of probability measures, follow from radial and angular distribution of Galois conjugates of algebraic numbers with small height. Here, radial distribution means that “most of” these conjugates have absolute value close to 1. It is easily established from the definition of the Weil height. The angular distribution deals with the distribution in $[0, 2\pi]$ of arguments of these conjugates modulo 2π . It is the hardest part of Mignotte’s proof. The first idea is to apply a result of Erdős and Turán [16] (see Theorem 2.3 in Section 2) which asserts that the arguments of roots of a polynomial $Q(X) = \sum_{i=0}^D q_i X^i$ with complex coefficients are well distributed in $[0, 2\pi]$ if the ratio

$$\frac{\mathcal{L}(Q)}{\sqrt{|q_D q_0|}} \tag{1.1}$$

is “not too large”. Here, $\mathcal{L}(Q) = \sum_{i=0}^D |q_i|$ denotes the length of Q . Unfortunately, the coefficients of the minimal polynomial of an algebraic number α can be very large, even if the height of α is small. This is when the second ingredient of the proof comes in. A classical result in diophantine approximation, the Siegel Lemma, shows that there exists a polynomial $Q \in \mathbb{Z}[X]$ of “small” degree vanishing at α such that $q_0 q_D$ is non-zero and its coefficients are “not too large” if the height of α is small. The quantity in (1.1) is therefore “not too large” since $|q_0 q_D| \geq 1$.

In the nineties, Mignotte pointed out to the first author that to obtain the angular distribution, we do not need to have an auxiliary polynomial with integer coefficients, but only with $|q_0 q_D|$ not too small. This innocent remark was one of the starting points of our investigation.

In this article, we apply Mignotte’s method to deal with the radial and angular distribution of sets S with large cardinality and made of algebraic numbers with small height. The novelty is that our sets are no more assumed to be stable by Galois action. This prevents us from using the standard Siegel Lemma, which makes the study of the angular distribution more complicated. Fortunately, there is an absolute version of this lemma, which follows from deep results of Zhang [29, Theorem 1.10]. This *absolute Siegel Lemma* can be applied to our situation, but it provides us an auxiliary polynomial $Q \in \overline{\mathbb{Q}}[X]$ whose coefficients (which can be assumed to be algebraic integers) cannot be controlled. In particular, $|q_0 q_D|$ can be as small as possible, but its absolute norm has to be a positive integer. This naturally leads us to consider the *arithmetic mean* of the radial and angular discrepancy of the conjugate sets of S . We prove that both of them are small.

Let $r > 1$ be a real. We write \mathcal{A}_r for the closed annulus centred at the origin with inner radius $1/r$ and outer radius r . We also denote by $\bar{h}(S)$ the (arithmetic) mean of $h(\alpha)$ with $\alpha \in S$, that is,

$$\bar{h}(S) = \frac{1}{|S|} \sum_{\alpha \in S} h(\alpha).$$

We finally set

$$\bar{h}^*(S) = 24 \left(\bar{h}(S) + \frac{\log(2|S|)}{|S|} \right)^{1/3}.$$

Theorem 1.2. *Let $S \subset \overline{\mathbb{Q}}^*$ be a finite subset.*

(1) *For any $r > 1$, we have*

$$\frac{1}{[\mathbb{Q}(S) : \mathbb{Q}]} \sum_{\sigma: \mathbb{Q}(S) \rightarrow \mathbb{C}} \frac{|\sigma(S) \setminus \mathcal{A}_r|}{|S|} \leq \frac{2\bar{h}(S)}{\log r}.$$

(2) *For any sector Δ of angle $\theta \in [0, 2\pi]$ based at the origin, we have*

$$\frac{1}{[\mathbb{Q}(S) : \mathbb{Q}]} \sum_{\sigma: \mathbb{Q}(S) \rightarrow \mathbb{C}} \left| \frac{|\sigma(S) \cap \Delta|}{|S|} - \frac{\theta}{2\pi} \right| \leq \bar{h}^*(S).$$

When S is Galois invariant, we recover Mignotte's results [22]. The following example is a good illustration of what happens when we drop the assumption of Galois invariance on S . Let p be a prime number, and let $\zeta_p = \exp(2i\pi/p)$. We choose $S = S_p$ as the set of p -roots of unity ζ_p^k with $1 \leq k \leq \lfloor \sqrt{p} \rfloor$, where $[x]$ denotes as usual the integer part of $x \in \mathbb{R}$. Thus

$$\bar{h}^*(S_p) = 24 \left(\frac{\log(2\lfloor \sqrt{p} \rfloor)}{\lfloor \sqrt{p} \rfloor} \right)^{1/3} \rightarrow 0$$

as $p \rightarrow \infty$. We fix a sector Δ of angle $\theta \in [0, 2\pi]$ based at the origin with $1 \notin \Delta$. Then $S_p \cap \Delta$ is empty when p is sufficiently large; whence

$$\left| \frac{|S_p \cap \Delta|}{|S_p|} - \frac{\theta}{2\pi} \right| = \frac{\theta}{2\pi}.$$

However, by Theorem 1.2(2) (and as it can be directly verified),

$$\frac{1}{p-1} \sum_{\sigma: \mathbb{Q}(\zeta_p) \hookrightarrow \mathbb{C}} \left| \frac{|\sigma(S_p) \cap \Delta|}{|S_p|} - \frac{\theta}{2\pi} \right| \rightarrow 0$$

as $p \rightarrow \infty$. This means that there are a "small" number of \mathbb{Q} -embeddings $\sigma: \mathbb{Q}(\zeta_p) \rightarrow \mathbb{C}$ for which the ratio $|\sigma(S_p) \cap \Delta|/|S_p|$ is "far" from $\theta/(2\pi)$.

Theorem 1.2 allows us to prove new results on points of small Weil height. Following Bombieri and Zannier [10], we say that a set S of algebraic numbers has the *Bogomolov property*, or short property (B) , if there exists a positive constant c such that the Weil height of an element in S is either 0 or bounded from below by c . Property (B) was established for the maximal totally real extension \mathbb{Q}^{tr} of \mathbb{Q} by Schinzel, see [26]. Note that it is also an immediate consequence of Theorem 1.1. The following corollary can be viewed as a generalization of this result.

Corollary 1.3. *Let L be an algebraic field. Then for any α in the group product $L^*(\mathbb{Q}^{\text{tr}})^*$ such that $h(\alpha) < 5 \cdot 10^{-6}$, we have $[L(\alpha) : L] < 4 \cdot 10^6$.*

Proof. Assume by contradiction that there exists $\alpha \in L^*(\mathbb{Q}^{\text{tr}})^*$ such that $h(\alpha) < 5 \cdot 10^{-6}$ and $d = [L(\alpha) : L] \geq 4 \cdot 10^6$. We write $\alpha = yz$ with $y \in L^*$ and $z \in (\mathbb{Q}^{\text{tr}})^*$. Define S as the Galois orbit of α over L . Thus, for any \mathbb{Q} -embedding $\sigma: \mathbb{Q}(S) \rightarrow \mathbb{C}$, we have $\sigma(S) \subseteq \sigma(y)\mathbb{R}$. Hence, for any sector Δ based at the origin,

$$\frac{|\sigma(S) \cap \Delta|}{|S|} = \begin{cases} 1 & \text{if } \sigma(y) \in \Delta; \\ 0 & \text{otherwise.} \end{cases}$$

Thus, choosing for Δ any sector of angle π , the value of the left-hand side in Theorem 1.2(2) is $1/2$. On the other hand, $|S| = d$ and $\log(2d)/d < 4 \cdot 10^{-6}$ since $d \geq 4 \cdot 10^6$ by assumption. Moreover, $\bar{h}(S) = h(\alpha) < 5 \cdot 10^{-6}$. In conclusion, $\bar{h}^*(S) = 24(\bar{h}(S) + \log(2|S|)/|S|)^{1/3} < 1/2$, a contradiction. \square

The maximal abelian extension \mathbb{Q}^{ab} of \mathbb{Q} also satisfies property (B), as it has been conjectured by Zannier and proved in [2]. However, the compositum of \mathbb{Q}^{tr} and \mathbb{Q}^{ab} does not satisfy property (B) since its subfield $\mathbb{Q}^{\text{tr}}(i)$ does not satisfy (B), see [1, Theorem 5.3]. Nevertheless,

Corollary 1.4. *The group product $G = (\mathbb{Q}^{\text{ab}})^*(\mathbb{Q}^{\text{tr}})^*$ satisfies property (B).*

Proof. Assume by contradiction that there exists a sequence $(\alpha_n)_n$ belonging to $G \setminus \mu_\infty$ such that $h(\alpha_n) \rightarrow 0$. By [4, Theorem 1.1], $[\mathbb{Q}^{\text{ab}}(\alpha_n) : \mathbb{Q}^{\text{ab}}] \rightarrow +\infty$. We then apply Corollary 1.3 with $L = \mathbb{Q}^{\text{ab}}$. \square

Theorem 1.2 can be formulated in terms of convergence in distribution, as it was expected in [21] (see the paragraph therein around equation (15.9)). Let $r > 1$, and let $f : \mathbb{C}^* \rightarrow \mathbb{C}$ be a function that is Lipschitz on \mathcal{A}_r . We define $\text{Lip}_r(f)$ to be the Lipschitz constant of f on \mathcal{A}_r . The infinite norm of f on a set $T \subset \mathbb{C}^*$ is denoted with $\|f\|_{\infty, T}$. Finally, given a subset $S \subset \overline{\mathbb{Q}}^*$, we define $\overline{S}^{\text{Gal}}$ as the smallest $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -invariant set containing S .

Theorem 1.5. *Let $S \subset \overline{\mathbb{Q}}^*$ be a finite set, let $r > 1$ be a real number, and let $N \geq 2$ be an integer.*

(1) *For all functions $f : \mathbb{C}^* \rightarrow \mathbb{C}$ that are Lipschitz on \mathcal{A}_r , we have*

$$\begin{aligned} & \frac{1}{[\mathbb{Q}(S) : \mathbb{Q}]} \sum_{\sigma : \mathbb{Q}(S) \hookrightarrow \mathbb{C}} \left| \int_{\mathbb{C}^*} f d\mu_{\sigma(S)} - \int_{\mathbb{C}^*} f d\lambda \right| \\ & \leq \frac{4r\pi \text{Lip}_r(f)}{N} + (\|f\|_{\infty, \overline{S}^{\text{Gal}} \setminus \mathcal{A}_r} + 2\|f\|_{\infty, \mathcal{A}_r}) \frac{2\overline{h}(S)}{\log r} + 2N\|f\|_{\infty, \mathcal{A}_r} \overline{h}^*(S). \end{aligned}$$

(2) *Let $\varepsilon \in (0, 1)$, and let L be a number field. Then there exists a set $\Lambda = \Lambda(S, r, N, \varepsilon, L)$ of L -embeddings $L(S) \hookrightarrow \mathbb{C}$ with cardinality at least $(1 - \varepsilon)[L(S) : L]$ such that*

$$\begin{aligned} & \left| \int_{\mathbb{C}^*} f d\mu_{\sigma(S)} - \int_{\mathbb{C}^*} f d\lambda \right| \leq \frac{4r\pi \text{Lip}_r(f)}{N} \\ & + (\|f\|_{\infty, \overline{S}^{\text{Gal}} \setminus \mathcal{A}_r} + 2\|f\|_{\infty, \mathcal{A}_r}) \frac{4[L : \mathbb{Q}]\overline{h}(S)}{\varepsilon \log r} + \frac{4[L : \mathbb{Q}]N^2}{\varepsilon} \|f\|_{\infty, \mathcal{A}_r} \overline{h}^*(S) \end{aligned}$$

for all $\sigma \in \Lambda$ and all functions $f : \mathbb{C}^ \rightarrow \mathbb{C}$ that are Lipschitz on \mathcal{A}_r .*

Theorem 1.5 implies a quantitative version of Bilu's equidistribution theorem. Such versions already exist in the literature, see [23, 17, 13, 6]. In all these articles, the estimations are stronger than ours, but they can only hold for much more restrictive functions. For instance, in the first three references, f must be at least bounded and differentiable on \mathbb{C}^* and in the last one, f has to be Lipschitz on \mathcal{A}_r , continuous on \mathbb{C}^* and satisfy $|f(z)| \leq \log |z|$

when $|z| \geq r$ as well as $|f(z)| \leq \log |z|^{-1}$ when $0 < |z| \leq 1/r$. Regarding the test functions of Theorem 1.5, they must be Lipschitz on \mathcal{A}_r , but can be unbounded and totally discontinuous outside.

Theorem 1.5 has the following qualitative consequence:

Corollary 1.6. *Let $(S_n)_n$ be a sequence of finite subsets of $\overline{\mathbb{Q}}^*$ such that $|S_n| \rightarrow +\infty$ and $\bar{h}(S_n) \rightarrow 0$. Let $V \subset \mathbb{C}^*$ be a neighbourhood of the unit circle. We consider the class of test functions $f : \mathbb{C}^* \rightarrow \mathbb{C}$ satisfying*

$$f \text{ is continuous on } V \quad \text{and} \quad \bar{h}(S_n) \|f\|_{\infty, S_n^{\text{Gal}} \setminus V} \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (1.2)$$

Then

(1) *We have*

$$\frac{1}{[\mathbb{Q}(S_n) : \mathbb{Q}]} \sum_{\sigma : \mathbb{Q}(S_n) \hookrightarrow \mathbb{C}} \left| \int_{\mathbb{C}^*} f d\mu_{\sigma(S_n)} - \int_{\mathbb{C}^*} f d\lambda \right| \rightarrow 0$$

for all functions $f : \mathbb{C}^* \rightarrow \mathbb{C}$ satisfying (1.2).

(2) *Let $\varepsilon \in (0, 1)$, and let L be a number field. Then for all integers $n \geq 1$, there is a set Λ_n of L -embeddings $L(S_n) \hookrightarrow \mathbb{C}$ with cardinality at least $(1 - \varepsilon)[L(S_n) : L]$ such that for all $(\sigma_n)_n \in \prod_n \Lambda_n$, we have*

$$\frac{1}{|S_n|} \sum_{\beta \in \sigma_n S_n} f(\beta) \rightarrow \int_0^1 f(e^{2i\pi t}) dt$$

for all functions $f : \mathbb{C}^* \rightarrow \mathbb{C}$ satisfying (1.2).

Note that every bounded continuous function $f : \mathbb{C}^* \rightarrow \mathbb{C}$ satisfies (1.2). Thus, the second assertion of this theorem implies:

Corollary 1.7. *Let $(S_n)_n$ be a sequence of finite subsets of $\overline{\mathbb{Q}}^*$ such that $|S_n| \rightarrow +\infty$ and $\bar{h}(S_n) \rightarrow 0$. Let L be a number field. Then for all integers $n \geq 1$, there is a set Λ_n of L -embeddings $L(S_n) \hookrightarrow \mathbb{C}$ with cardinality at least $(1 - \varepsilon)[L(S_n) : L]$ such that for all $(\sigma_n)_n \in \prod_n \Lambda_n$, the sequence of discrete probability measures $\mu_{\sigma_n S_n}$ converges in distribution to λ .*

Corollary 1.7 partially follows from [15, Theorem 3.17]. More precisely, under the same assumptions on S_n (which corresponds to a very special case of [15, Theorem 3.17]), Doyle, Fili and Tobin obtained the same conclusion than ours, but for the weak convergence. Their proof is based on potential theory, which is the other classical approach to deal with equidistribution.

Theorem 1.1 does not hold anymore if we relax the assumption “ K is a number field” to “ K is an algebraic field”. Indeed, put $\alpha_n = (1 - e^{2i\pi/n^n})^{1/n}$. Let S_n be the Galois orbit of α_n over $\mathbb{Q}(\mu_\infty)$. Clearly, $h(\alpha_n) \leq (\log 2)/n$ and $\alpha_n \notin \mu_\infty$ for all integers $n \geq 1$. Hence, $h(\alpha_n) \rightarrow 0$ and $|S_n| \rightarrow +\infty$.

Moreover, each element of S_n has absolute value $|\alpha_n|$. The series expansion of the exponential implies $\alpha_n \rightarrow 0$, and so the sequence μ_{S_n} cannot converge in distribution to λ . This elementary example shows that the conclusion of the next corollary is somehow optimal.

Corollary 1.8. *Let K be an algebraic field, and let $(\alpha_n)_n$ be a sequence of $\overline{\mathbb{Q}}^*$ such that $h(\alpha_n) \rightarrow 0$ and $[K(\alpha_n) : K] \rightarrow +\infty$. Let L be a number field. Then there is a set Λ_n of L -embeddings $L(S_n) \hookrightarrow \mathbb{C}$ with cardinality at least $(1 - \varepsilon)[L(S_n) : L]$ such that for all $(\sigma_n)_n \in \prod_n \Lambda_n$, the sequence of discrete probability measures $\mu_{\sigma_n S_n}$ converges in distribution to λ , where S_n is the Galois orbit of α_n over K .*

Proof. Take for S_n the set of conjugates of α_n over K in Corollary 1.7. \square

Plan of the article

The plan of the article is as follows. In Section 2, we implement our generalisation of Mignotte’s method to treat the radial and angular distributions, then we prove Theorem 1.2. In Section 3, we deduce Theorem 1.5 from Theorem 1.2 and Corollary 1.6 from Theorem 1.5.

Acknowledgement

We thank Fili, Sombra and Weiss for their fruitful discussions. The second author sincerely thanks the *Laboratoire de Mathématiques N. Oresme* of *Université de Caen Normandie* for housing him on May 2023 so that he can work face-to-face with the first author. He was also funded by the Morningside Center of Mathematics, CAS.

2 Radial and angular distribution

The proof of Theorem 1.2 is based on arguments due to Mignotte, which are well highlighted in [21, Chapter 15]. We will follow the exposition of this book. Throughout this section r denotes a real number greater than 1 and $S \subset \overline{\mathbb{Q}}^*$ a finite set.

2.1 Radial distribution

The goal of this subsection is to establish that the mean of the number of elements belonging to $\sigma(S) \setminus \mathcal{A}_r$, where σ ranges over all \mathbb{Q} -embeddings $\mathbb{Q}(S) \hookrightarrow \mathbb{C}$, cannot be “too large” if $\bar{h}(S)$ is small. Recall that δ_x is the Dirac measure on \mathbb{C}^* supported at x . Let us start by the following lemma.

Lemma 2.1 ([21], Theorem 15.1). *For all $\alpha \in \overline{\mathbb{Q}}^*$, we have*

$$\sum_{\tau: \mathbb{Q}(\alpha) \hookrightarrow \mathbb{C}} \delta_{\tau\alpha}(\mathcal{A}_r) \geq [\mathbb{Q}(\alpha) : \mathbb{Q}] \left(1 - \frac{2h(\alpha)}{\log r} \right).$$

We can now prove the “radial part” of Theorem 1.2.

Proof of Theorem 1.2(1). Since $|\sigma(S) \cap \mathcal{A}_r| + |\sigma(S) \setminus \mathcal{A}_r| = |S|$, the inequality in Theorem 1.2(1) is equivalent to

$$\frac{1}{[\mathbb{Q}(S) : \mathbb{Q}]} \sum_{\sigma: \mathbb{Q}(S) \hookrightarrow \mathbb{C}} |\sigma(S) \cap \mathcal{A}_r| \geq |S| \left(1 - \frac{2\bar{h}(S)}{\log r}\right),$$

which we now prove. The set $\sigma(S) \cap \mathcal{A}_r$ has cardinality $\sum_{\alpha \in S} \delta_{\sigma\alpha}(\mathcal{A}_r)$ for all $\sigma: \mathbb{Q}(S) \hookrightarrow \mathbb{C}$. Thus,

$$\sum_{\sigma: \mathbb{Q}(S) \hookrightarrow \mathbb{C}} |\sigma(S) \cap \mathcal{A}_r| = \sum_{\alpha \in S} \sum_{\sigma: \mathbb{Q}(S) \hookrightarrow \mathbb{C}} \delta_{\sigma\alpha}(\mathcal{A}_r).$$

Then, each \mathbb{Q} -embedding $\tau: \mathbb{Q}(\alpha) \hookrightarrow \mathbb{C}$ can be extended in $[\mathbb{Q}(S) : \mathbb{Q}(\alpha)]$ different ways to a \mathbb{Q} -embedding from $\mathbb{Q}(S)$ to \mathbb{C} , which leads to

$$\sum_{\sigma: \mathbb{Q}(S) \hookrightarrow \mathbb{C}} |\sigma(S) \cap \mathcal{A}_r| = \sum_{\alpha \in S} [\mathbb{Q}(S) : \mathbb{Q}(\alpha)] \sum_{\tau: \mathbb{Q}(\alpha) \hookrightarrow \mathbb{C}} \delta_{\tau\alpha}(\mathcal{A}_r).$$

By Lemma 2.1, the right-hand side is at least $[\mathbb{Q}(S) : \mathbb{Q}] \sum_{\alpha \in S} \left(1 - \frac{2h(\alpha)}{\log r}\right)$. Theorem 1.2(1) now arises from a small calculation. \square

2.2 Angular distribution

This subsection aims to show that the elements of $\sigma(S)$ are, in average, angularly well distributed when $\bar{h}^*(S)$ is small. Concretely, if Δ is a sector of angle θ based at the origin, then the mean of number of elements belonging to $\sigma(S) \cap \Delta$, where σ runs over all \mathbb{Q} -embeddings $\mathbb{Q}(S) \rightarrow \mathbb{C}$, is approximately $\theta|S|/(2\pi)$ when $\bar{h}^*(S)$ is small.

Remark 2.2. *The left-hand side in Theorem 1.2(2) is obviously bounded from above by 1. Hence, the theorem is trivial unless $\bar{h}^*(S) \leq 1$. Moreover, if $\bar{h}^*(S) \leq 1$, then the definition of $\bar{h}^*(S)$ implies $24(\log(2|S|)/|S|)^{1/3} \leq 1$, and so $|S| \geq 3$. To summarize, we can reduce the proof of Theorem 1.2(2) to the case that $\bar{h}^*(S) \leq 1$ and $|S| \geq 3$, what we now assume.*

The proof of Theorem 1.2(2) is mainly based on two ingredients. The first one is a result due to Erdős and Turán [16], see also [3] for a more modern proof of something slightly sharper.

For any region $\Delta \subset \mathbb{C}$ and any polynomial $Q \in \mathbb{C}[X]$, we denote by $Z_{\Delta, Q}$ the number of zeroes of Q (with multiplicity) lying in Δ .

Theorem 2.3 (Erdős-Turán). *Let Δ be a sector of angle $\theta \in [0, 2\pi]$ based at the origin, and let $Q(X) = \sum_{i=0}^D q_i X^i \in \mathbb{C}[X]$ be a polynomial with $q_D q_0 \neq 0$. Then*

$$\left| Z_{\Delta, Q} - \frac{\theta}{2\pi} D \right|^2 \leq 256D \log \left(\frac{\mathcal{L}(Q)}{\sqrt{|q_D q_0|}} \right),$$

where $\mathcal{L}(Q) = \sum_{i=0}^D |q_i|$ denotes the length of Q .

Write $P_S \in \mathbb{Q}(S)[X]$ for the polynomial $\prod_{\alpha \in S} (X - \alpha)$. Thus $|\sigma(S) \cap \Delta| = Z_{\Delta, \sigma P_S}$. The natural idea to get Theorem 1.2(2) would be to apply Theorem 2.3 to $Q = \sigma P_S$ with σ running over all \mathbb{Q} -embeddings $\mathbb{Q}(S) \hookrightarrow \mathbb{C}$. But the mean of $\log(\mathcal{L}(\sigma P_S))$ might be too large, spoiling our chances of getting what we wish. This is when the second ingredient comes in: the absolute Siegel's lemma.

Let $F(X) = \sum_{i=0}^L f_i X^i \in \overline{\mathbb{Q}}[X]$ be a polynomial. The height of F , denoted with $h(F)$, is the Weil height of coefficients of F , that is,

$$h(F) = \frac{1}{[E : \mathbb{Q}]} \sum_v [E_v : \mathbb{Q}_v] \log \max\{|f_0|_v, \dots, |f_L|_v\},$$

where E is any number field containing f_0, \dots, f_L and where v ranges over all places of E . It is well-known that this definition does not depend on the choice of such a field E .

Theorem 2.4 (Absolute Siegel's Lemma). *Let L be a positive integer with $L > |S|$. Then there exists a non-zero polynomial F , with algebraic integer coefficients and degree $< L$, vanishing at S such that:*

$$h(F) \leq \frac{|S|}{L+1-|S|} \left(\frac{3}{2} \log(L+1) + (L+1)\overline{h}(S) \right) + \frac{\log(L+1)}{2}.$$

This statement improves the main result of Roy and Thunder, see [24, Theorem 2.2]. It is an easy consequence of [29, Theorem 5.2], see [4, Proposition 4.2] for details¹.

Define L as the round up to $(1 + \overline{h}^*(S)/6)|S|$. By Remark 2.2, we have $L \leq 2|S| - 1$. By Theorem 2.4 and using the inequalities

$$\left(1 + \frac{\overline{h}^*(S)}{6} \right) |S| \leq L + 1 \leq 2|S|,$$

we find a non-zero polynomial $F(X) = \sum_{i=0}^L f_i X^i$, with algebraic integer coefficients and degree $< L$, divisible by P_S such that

$$h(F) \leq \frac{6}{\overline{h}^*(S)} \left(\frac{3}{2} \log(2|S|) + 2|S|\overline{h}(S) \right) + \frac{\log(2|S|)}{2}. \quad (2.1)$$

¹In *op.cit.* the relevant height is $h_2(F)$, with the L_2 -metric at archimedean places, but obviously $h(F) \leq h_2(F)$.

Dividing F by a power of X if needed, we can assume that $f_D f_0 \neq 0$. We now choose a number field E containing $\mathbb{Q}(S)$ and all coefficients of F .

To prove Theorem 1.1 via this approach (see [21, Chapter 15]), S is the set of all Galois conjugates of some $\alpha \in \overline{\mathbb{Q}}^*$. Thus, P_S is the minimal polynomial of α over \mathbb{Q} and the classical Siegel's lemma asserts that we can take F with integer coefficients, which implies $|f_D f_0| \geq 1$. Unfortunately, in our situation, we can have $|f_D f_0| \neq 0$ as small as possible. Nonetheless, $f_D f_0$ is an algebraic integer and therefore its norm $\prod_{\sigma: E \hookrightarrow \mathbb{C}} |\sigma(f_D f_0)|$ over \mathbb{Q} is a positive integer.

Lemma 2.5. *Let Δ be a sector of angle $\theta \in [0, 2\pi]$ based at the origin. Then*

$$\frac{1}{[E : \mathbb{Q}]} \sum_{\sigma: E \hookrightarrow \mathbb{C}} \left| Z_{\Delta, \sigma F} - \frac{\theta}{2\pi} D \right|^2 \leq \left(\frac{2|S|\bar{h}^*(S)}{3} \right)^2.$$

Proof. Let $\sigma: E \hookrightarrow \mathbb{C}$ be a \mathbb{Q} -embedding. Then,

$$\mathcal{L}(\sigma F) = \sum_{i=0}^D |\sigma f_i| \leq (D+1) \max\{|\sigma f_0|, \dots, |\sigma f_D|\}.$$

By (2.1), we get

$$\begin{aligned} \frac{1}{[E : \mathbb{Q}]} \sum_{\sigma: E \hookrightarrow \mathbb{C}} \log(\mathcal{L}(\sigma F)) &\leq \log(D+1) + h(F) \\ &\leq \frac{3}{2} \log(2|S|) + \frac{6}{\bar{h}^*(S)} \left(\frac{3}{2} \log(2|S|) + 2|S|\bar{h}(S) \right) \\ &\leq \frac{12}{\bar{h}^*(S)} (\log(2|S|) + |S|\bar{h}(S)) \end{aligned}$$

because $D < L \leq 2|S| - 1$ and $\bar{h}^*(S) \leq 1$. By definition of $\bar{h}^*(S)$, we conclude

$$\frac{1}{[E : \mathbb{Q}]} \sum_{\sigma: E \hookrightarrow \mathbb{C}} \log(\mathcal{L}(\sigma F)) \leq \frac{12|S|}{\bar{h}^*(S)} \left(\frac{\bar{h}^*(S)}{24} \right)^3 = \frac{|S|\bar{h}^*(S)^2}{2^7 \cdot 3^2}.$$

Finally, Theorem 2.3 applied to $Q = \sigma F$ gives

$$\sum_{\sigma: E \hookrightarrow \mathbb{C}} \left| Z_{\Delta, \sigma F} - \frac{\theta}{2\pi} D \right|^2 \leq 2^9 |S| \sum_{\sigma: E \hookrightarrow \mathbb{C}} \log(\mathcal{L}(\sigma F))$$

since $\sum_{\sigma} \log |\sigma(f_D f_0)| \geq 0$ by the foregoing. The lemma follows. \square

Proof of Theorem 1.2(2). Note that the left-hand side in Theorem 1.2(2) remains unchanged if we replace $\mathbb{Q}(S)$ with a finite extension. So, it is

enough to prove Theorem 1.2(2) by replacing $\mathbb{Q}(S)$ with E . Let $\sigma: E \hookrightarrow \mathbb{C}$ be a \mathbb{Q} -embedding. The triangle inequality gives

$$\left| Z_{\Delta, \sigma P_S} - \frac{\theta |S|}{2\pi} \right| \leq |Z_{\Delta, \sigma P_S} - Z_{\Delta, \sigma F}| + \left| Z_{\Delta, \sigma F} - \frac{\theta D}{2\pi} \right| + \left| \frac{\theta D}{2\pi} - \frac{\theta |S|}{2\pi} \right|.$$

Recall that P_S divides F . Thus, $Z_{\Delta, \sigma F} - Z_{\Delta, \sigma P_S} = Z_{\Delta, \sigma(F/P_S)}$. In particular, it is bounded from above by the degree of F/P_S , namely $D - |S|$, and so by $L - 1 - |S| \leq |S| \bar{h}^*(S)/6$. Thus,

$$\frac{1}{[E : \mathbb{Q}]} \sum_{\sigma: E \hookrightarrow \mathbb{C}} \left| Z_{\Delta, \sigma P_S} - \frac{\theta}{2\pi} |S| \right| \leq \frac{|S| \bar{h}^*(S)}{3} + \frac{1}{[E : \mathbb{Q}]} \sum_{\sigma: E \hookrightarrow \mathbb{C}} \left| Z_{\Delta, \sigma F} - \frac{\theta}{2\pi} D \right|.$$

From Lemma 2.5, we have

$$\begin{aligned} \frac{1}{[E : \mathbb{Q}]} \sum_{\sigma: E \hookrightarrow \mathbb{C}} \left| Z_{\Delta, \sigma F} - \frac{\theta}{2\pi} D \right| &\leq \sqrt{\frac{1}{[E : \mathbb{Q}]} \sum_{\sigma: E \hookrightarrow \mathbb{C}} \left| Z_{\Delta, \sigma F} - \frac{\theta}{2\pi} D \right|^2} \\ &\leq \frac{2|S| \bar{h}^*(S)}{3} \end{aligned}$$

and the second assertion of Theorem 1.2 follows. \square

3 Convergence in distribution

Let S and r be as in Theorem 1.5. By Theorem 1.2, we obtain that in average, the cardinality of $\sigma(S) \cap \Delta \cap \mathcal{A}_r$, with $\sigma: \mathbb{Q}(S) \hookrightarrow \mathbb{C}$ a \mathbb{Q} -embedding, is approximately $\theta|S|/(2\pi)$ when $\bar{h}(S)$ and $\bar{h}^*(S)$ are small enough. This assertion is stronger when θ is small. For this reason, it makes sense to cut \mathcal{A}_r into a large number of small annulus sectors, apply Theorem 1.2 to deduce that in average, there are around $\theta|S|/(2\pi)$ elements of $\sigma(S)$ in each of these annulus sectors, and put it all the information together to conclude that in average, the set $\sigma(S)$ is equidistributed around the unit circle. Formalizing this process leads to the proof of Theorem 1.5(1). A slight modification of these arguments shows the second part.

We start with an easy lemma. Given $r > 1$ and $t_0, t_1 \in [0, 1]$ with $t_0 < t_1$, we consider the compact region

$$V_{r, t_0, t_1} = \{\rho e^{2i\pi t} \mid \rho \in [1/r, r], t \in [t_0, t_1]\}.$$

Lemma 3.1. *Let $f: \mathbb{C}^* \rightarrow \mathbb{R}$ be a real function that is Lipschitz on \mathcal{A}_r for some $r > 1$, and let T be a finite set of non-zero complex numbers. Choose $t_0, t_1 \in [0, 1]$ with $t_0 < t_1$ and put $V = V_{r, t_0, t_1}$. If $\theta = t_1 - t_0 \in (0, 1/2]$, then*

$$\left| \frac{1}{|T|} \sum_{\beta \in T \cap V} f(\beta) - \int_{t_0}^{t_1} f(e^{2i\pi t}) dt \right| \leq 2r\pi\theta^2 \text{Lip}_r(f) + \|f\|_{\infty, \mathcal{A}_r} \left| \frac{|T \cap V|}{|T|} - \theta \right|.$$

Proof. Let f^+ , resp. f^- , be the supremum, resp. infimum, of $f(z)$, where z ranges over all elements of V . Then

$$\frac{f^-|T \cap V|}{|T|} - f^+\theta \leq \frac{1}{|T|} \sum_{\beta \in T \cap V} f(\beta) - \int_{t_0}^{t_1} f(e^{2i\pi t}) dt \leq \frac{f^+|T \cap V|}{|T|} - f^-\theta.$$

We write

$$\frac{f^\pm|T \cap V|}{|T|} - f^\mp\theta = f^\pm \left(\frac{|T \cap V|}{|T|} - \theta \right) + (f^\pm - f^\mp)\theta.$$

Combining the last two displayed equations, then using the triangle inequality, we get

$$\left| \frac{1}{|T|} \sum_{\beta \in T \cap V} f(\beta) - \int_{t_0}^{t_1} f(e^{2i\pi t}) dt \right| \leq (f^+ - f^-)\theta + \|f\|_{\infty, \mathcal{A}_r} \left| \frac{|T \cap V|}{|T|} - \theta \right|. \quad (3.1)$$

The fact that V is compact implies that $f^+ = f(x)$ and $f^- = f(y)$ for some $x, y \in V$. Each point $z \in V$ expresses as $r_z e^{i\alpha_z}$ with $r_z \in [1/r, r]$ and $\alpha_z \in [2\pi t_0, 2\pi t_1]$. Hence,

$$\begin{aligned} f^+ - f^- &= |f(x) - f(y)| \leq \text{Lip}_r(f) |r_x e^{i\alpha_x} - r_y e^{i\alpha_y}| \\ &= \text{Lip}_r(f) \sqrt{r_x^2 + r_y^2 - 2r_x r_y \cos(\alpha_y - \alpha_x)}. \end{aligned}$$

As $\theta \in (0, 1/2]$, the cosine is decreasing on $[0, 2\pi\theta]$. Thus $\cos(\alpha_y - \alpha_x) \geq \cos(2\pi\theta)$, and so

$$|f^+ - f^-| \leq \text{Lip}_r(f) \sqrt{2r^2(1 - \cos(2\pi\theta))} \quad (3.2)$$

since $r_x, r_y \leq r$. The lemma follows by combining (3.1) and (3.2), then using the inequality $1 - \cos(x) \leq x^2/2$, which is true for all $x \in \mathbb{R}$. \square

Fix from now a finite subset $S \subset \overline{\mathbb{Q}}^*$ as well as an integer $N \geq 2$. For $x \in \mathbb{R}$ and $j \in \{0, \dots, N-1\}$, we denote by $\Delta_j(x)$ the sector based at the origin containing 0 and all non-zero complex numbers whose argument belongs to $[x+2\pi j/N, x+2\pi(j+1)/N]$ up to $2k\pi$. It is a sector of angle $2\pi/N$. The Galois closure $\overline{S}^{\text{Gal}}$ of S being finite, we can then find $x \in \mathbb{R}$ satisfying the following property: for all $\alpha \in \overline{S}^{\text{Gal}}$, there is (a unique) $j \in \{0, \dots, N-1\}$ such that α lies in the interior of $\Delta_j(x)$. We fix from now such an x and in order to ease notation, we put $\Delta_j = \Delta_j(x)$.

Let $\sigma: \mathbb{Q}(S) \hookrightarrow \mathbb{C}$ be a \mathbb{Q} -embedding, and let $f: \mathbb{C}^* \rightarrow \mathbb{C}$ be a function that is Lipschitz on \mathcal{A}_r . We have

$$\begin{aligned} \left| \int_{\mathbb{C}^*} f d\mu_{\sigma(S)} - \int_{\mathbb{C}^*} f d\lambda \right| &= \left| \frac{1}{|S|} \sum_{\beta \in \sigma(S)} f(\beta) - \int_0^1 f(e^{2i\pi t}) dt \right| \\ &\leq \frac{|\sigma(S) \setminus \mathcal{A}_r|}{|S|} \|f\|_{\infty, \overline{S}^{\text{Gal}} \setminus \mathcal{A}_r} + \left| \frac{1}{|S|} \sum_{\beta \in \sigma(S) \cap \mathcal{A}_r} f(\beta) - \int_0^1 f(e^{2i\pi t}) dt \right|. \end{aligned} \quad (3.3)$$

As each element of $\sigma(S)$ lies in Δ_j for a unique $j \in \{0, \dots, N-1\}$, we get

$$\begin{aligned} \frac{1}{|S|} \sum_{\beta \in \sigma(S) \cap \mathcal{A}_r} f(\beta) - \int_0^1 f(e^{2i\pi t}) dt \\ = \sum_{j=0}^{N-1} \left(\frac{1}{|S|} \sum_{\beta \in \sigma(S) \cap \Delta_j \cap \mathcal{A}_r} f(\beta) - \int_{j/N}^{(j+1)/N} f(e^{2i\pi t}) dt \right). \end{aligned}$$

The real and imaginary parts of f are two real-valued functions which are Lipschitz on \mathcal{A}_r since f is. Moreover, their Lipschitz coefficients and suprema on \mathcal{A}_r are bounded from above by those of f . Applying Lemma 3.1 twice to the real part (for the first time), then the imaginary part (for the second one) and with $t_0 = j/N$, $t_1 = (j+1)/N$ and $T = \sigma(S)$ in both cases, we conclude thanks to the triangle inequality that

$$\begin{aligned} \left| \frac{1}{|S|} \sum_{\beta \in \sigma(S) \cap \Delta_j \cap \mathcal{A}_r} f(\beta) - \int_{j/N}^{(j+1)/N} f(e^{2i\pi t}) dt \right| \\ \leq \frac{4r\pi \text{Lip}_r(f)}{N^2} + 2\|f\|_{\infty, \mathcal{A}_r} \left| \frac{|\sigma(S) \cap \Delta_j \cap \mathcal{A}_r|}{|S|} - \frac{1}{N} \right|. \end{aligned}$$

We now infer that

$$\begin{aligned} \left| \frac{1}{|S|} \sum_{\beta \in \sigma(S) \cap \mathcal{A}_r} f(\beta) - \int_0^1 f(e^{2i\pi t}) dt \right| \\ \leq \frac{4r\pi \text{Lip}_r(f)}{N} + 2\|f\|_{\infty, \mathcal{A}_r} \sum_{j=0}^{N-1} \left| \frac{|\sigma(S) \cap \Delta_j \cap \mathcal{A}_r|}{|S|} - \frac{1}{N} \right|. \end{aligned} \quad (3.4)$$

Since $|\sigma(S) \cap \Delta_j \cap \mathcal{A}_r| = |\sigma(S) \cap \Delta_j| - |\sigma(S) \cap \Delta_j \setminus \mathcal{A}_r|$, the triangle inequality gives

$$\left| \frac{|\sigma(S) \cap \Delta_j \cap \mathcal{A}_r|}{|S|} - \frac{1}{N} \right| \leq \left| \frac{|\sigma(S) \cap \Delta_j|}{|S|} - \frac{1}{N} \right| + \frac{|\sigma(S) \cap \Delta_j \setminus \mathcal{A}_r|}{|S|}.$$

Since \mathbb{C} is the union of all $\Delta_0, \dots, \Delta_{N-1}$ and each element of $\sigma(S)$ lies in the interior of Δ_j for a unique $j \in \{0, \dots, N-1\}$, we get

$$\sum_{j=0}^{N-1} |\sigma(S) \cap \Delta_j \setminus \mathcal{A}_r| = |\sigma(S) \setminus \mathcal{A}_r|.$$

Thus,

$$\sum_{j=0}^{N-1} \left| \frac{|\sigma(S) \cap \Delta_j \cap \mathcal{A}_r|}{|S|} - \frac{1}{N} \right| \leq \frac{|\sigma(S) \setminus \mathcal{A}_r|}{|S|} + \sum_{j=0}^{N-1} \left| \frac{|\sigma(S) \cap \Delta_j|}{|S|} - \frac{1}{N} \right|. \quad (3.5)$$

By (3.3), (3.4) and (3.5), we get:

$$\begin{aligned} \left| \int_{\mathbb{C}^*} f d\mu_{\sigma(S)} - \int_{\mathbb{C}^*} f d\lambda \right| &\leq \frac{4r\pi \text{Lip}_r(f)}{N} + (\|f\|_{\infty, \overline{S}^{\text{Gal}} \setminus \mathcal{A}_r} + 2\|f\|_{\infty, \mathcal{A}_r}) \frac{|\sigma(S) \setminus \mathcal{A}_r|}{|S|} \\ &\quad + 2\|f\|_{\infty, \mathcal{A}_r} \sum_{j=0}^{N-1} \left| \frac{|\sigma(S) \cap \Delta_j|}{|S|} - \frac{1}{N} \right|. \quad (3.6) \end{aligned}$$

Proof of Theorem 1.5(1). It now arises from Theorem 1.2 (applied to the sector $\Delta = \Delta_j$ of angle $2\pi/N$) and from (3.6) that

$$\begin{aligned} &\frac{1}{[\mathbb{Q}(S) : \mathbb{Q}]} \sum_{\sigma: \mathbb{Q}(S) \hookrightarrow \mathbb{C}} \left| \int_{\mathbb{C}^*} f d\mu_{\sigma(S)} - \int_{\mathbb{C}^*} f d\lambda \right| \\ &\leq \frac{4r\pi \text{Lip}_r(f)}{N} + (\|f\|_{\infty, \overline{S}^{\text{Gal}} \setminus \mathcal{A}_r} + 2\|f\|_{\infty, \mathcal{A}_r}) \frac{2\overline{h}(S)}{\log r} + 2N\|f\|_{\infty, \mathcal{A}_r} \overline{h}^*(S), \end{aligned}$$

which is the first inequality of the theorem. \square

Proof of Theorem 1.5(2). We now prove the second assertion. Note that in the arithmetic means of Theorem 1.2, we may replace $\mathbb{Q}(S)$ with any of its finite field extensions. Thus, the first assertion of Theorem 1.2 easily implies that the set of \mathbb{Q} -embeddings $\sigma: L(S) \hookrightarrow \mathbb{C}$ satisfying

$$\frac{|\sigma(S) \setminus \mathcal{A}_r|}{|S|} \leq \frac{4[L : \mathbb{Q}] \overline{h}(S)}{\varepsilon \log r} \quad (3.7)$$

has cardinality at least $(1 - \frac{\varepsilon}{2[L:\mathbb{Q}]})[L(S) : \mathbb{Q}]$. Let $j \in \{0, \dots, N-1\}$. Similarly, the second assertion of Theorem 1.2 (with $\Delta = \Delta_j$) provides at least $(1 - \frac{\varepsilon}{2N[L:\mathbb{Q}]})[L(S) : \mathbb{Q}]$ field embeddings $\sigma: L(S) \hookrightarrow \mathbb{C}$ for which

$$\left| \frac{|\sigma(S) \cap \Delta_j|}{|S|} - \frac{1}{N} \right| \leq \frac{2N[L : \mathbb{Q}]}{\varepsilon} \overline{h}^*(S). \quad (3.8)$$

Thus, there exists a set $Y = Y(S, r, N, \varepsilon, L)$ of \mathbb{Q} -embeddings $\sigma : L(S) \hookrightarrow \mathbb{C}$ with cardinality

$$|Y| \geq \left(1 - \frac{\varepsilon}{[L : \mathbb{Q}]}\right) [L(S) : \mathbb{Q}] = [L(S) : \mathbb{Q}] - \varepsilon[L(S) : L]$$

such that any σ in this set satisfies (3.7) and (3.8) for all $j \in \{0, \dots, N-1\}$. We can moreover find a set $\Lambda \subseteq Y$ of L -embeddings with cardinality at least $(1 - \varepsilon)[L(S) : L]$. Otherwise, the cardinality of Y would be less than

$$(1 - \varepsilon)[L(S) : L] + ([L : \mathbb{Q}] - 1)[L(S) : L] = [L(S) : \mathbb{Q}] - \varepsilon[L(S) : L],$$

a contradiction. We choose such a subset Λ and we fix $\sigma \in \Lambda$.

Theorem 1.5(2) follows since by (3.6), (3.7) and (3.8), we get

$$\begin{aligned} \left| \int_{\mathbb{C}^*} f d\mu_{\sigma(S)} - \int_{\mathbb{C}^*} f d\lambda \right| &\leq \frac{4r\pi \text{Lip}_r(f)}{N} \\ &+ (\|f\|_{\infty, \overline{S}^{\text{Gal}} \setminus \mathcal{A}_r} + 2\|f\|_{\infty, \mathcal{A}_r}) \frac{4[L : \mathbb{Q}] \overline{h}(S)}{\varepsilon \log r} + \frac{4[L : \mathbb{Q}] N^2}{\varepsilon} \|f\|_{\infty, \mathcal{A}_r} \overline{h}^*(S) \end{aligned}$$

for all functions $f : \mathbb{C}^* \rightarrow \mathbb{C}$ that are Lipschitz on \mathcal{A}_r . \square

Proof of Corollary 1.6. We only prove the second assertion, the proof of the first one following similar lines. Given an integer n , a \mathbb{Q} -embedding $\sigma : \mathbb{Q}(S_n) \hookrightarrow \mathbb{C}$ and a function $f : \mathbb{C}^* \rightarrow \mathbb{C}$, we let for short

$$u_{n, \sigma}(f) = \left| \int_{\mathbb{C}^*} f d\mu_{\sigma S_n} - \int_{\mathbb{C}^*} f d\lambda \right|.$$

Let $f : \mathbb{C}^* \rightarrow \mathbb{C}$ be a function satisfying (1.2). Obviously, there exists $r > 1$ such that f is continuous on \mathcal{A}_r . Since f is not necessarily Lipschitz on \mathcal{A}_r , we use a standard density argument. Thanks to the Stone-Weierstrass theorem, we know that f restricted to \mathcal{A}_r is the uniform limit of a sequence of polynomial functions $(f_m)_m$. We extend f_m to a function on \mathbb{C}^* by setting $f_m = f$ on $\mathbb{C}^* \setminus \mathcal{A}_r$. Thus f_m is Lipschitz on \mathcal{A}_r for all m and $(f_m)_m$ uniformly converges to f on \mathbb{C}^* .

From our assumptions, we have $\overline{h}^*(S_n) \rightarrow 0$. The second assertion of Theorem 1.5 with $S = S_n$ and $N = \lceil \overline{h}^*(S_n)^{-1/4} \rceil$ proves that there is a set Λ_n of L -embeddings $\sigma : L(S_n) \hookrightarrow \mathbb{C}$, depending only on S_n, r, ε and L , but not on the test function f , such that $|\Lambda_n| \geq (1 - \varepsilon)[L(S_n) : L]$ and

$$\begin{aligned} u_{n, \sigma}(f_m) \leq U_{n, m} &= \frac{4r\pi \text{Lip}_r(f_m)}{\lceil \overline{h}^*(S_n)^{-1/4} \rceil} + (\|f_m\|_{\infty, S_n^{\text{Gal}} \setminus \mathcal{A}_r} + 2\|f_m\|_{\infty, \mathcal{A}_r}) \frac{4[L : \mathbb{Q}] \overline{h}(S_n)}{\varepsilon \log r} \\ &+ \frac{4[L : \mathbb{Q}] \lceil \overline{h}^*(S_n)^{-1/4} \rceil^2}{\varepsilon} \|f_m\|_{\infty, \mathcal{A}_r} \overline{h}^*(S_n) \quad (3.9) \end{aligned}$$

for all m . For each n , we choose one of these L -embeddings, say σ_n . We want to show that $u_{n,\sigma_n}(f) \rightarrow 0$, which would show the corollary.

Let m be an index. As $f = f_m$ outside \mathcal{A}_r , we deduce by assumption that $\bar{h}(S_n)\|f_m\|_{\infty, S_n^{\text{Gal}} \setminus \mathcal{A}_r} \rightarrow 0$, and so (3.9) leads to $U_{n,m} \rightarrow 0$ as $n \rightarrow +\infty$.

Let n be an index. Using the reverse triangle inequality, then the triangle inequality, we get

$$|u_{n,\sigma_n}(f_m) - u_{n,\sigma_n}(f)| \leq \int_{\mathbb{C}^*} |f_m - f| d\mu_{\sigma_n S_n} + \int_{\mathbb{C}^*} |f_m - f| d\lambda$$

for all m . As f_m uniformly converges to f on \mathbb{C}^* , we deduce that $u_{n,\sigma_n}(f_m)$ uniformly converges to $u_{n,\sigma_n}(f)$ as $m \rightarrow +\infty$. The Moore-Osgood theorem for interchanging limits leads to $u_{n,\sigma_n}(f) \rightarrow 0$ as $n \rightarrow +\infty$. \square

References

- [1] F. Amoroso, S. David and U. Zannier, *On fields with the property (B)*, Proc. Amer. Math. Soc. **142** (2014), p. 893–1910.
- [2] F. Amoroso and R. Dvornicich, *A Lower Bound for the Height in Abelian Extensions*, J. Number Theory **80** (2000), p. 260–272.
- [3] F. Amoroso and M. Mignotte, *On the distribution of the roots of polynomials*, Ann. Inst. Fourier (Grenoble) **46** (1996), p. 1275–1291.
- [4] F. Amoroso and U. Zannier, *A relative Dobrowolski lower bound over abelian extensions*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **29** (2000), p. 711–727.
- [5] M. Baker and L. Hsia, *Canonical heights, transfinite diameters, and polynomial dynamics*, J. Reine Angew. Math. **585** (2005), p. 61–92.
- [6] M. Baker and D. Masser, *Galois distribution on tori- A refinement, examples, and applications*, Int. Math. Res. Not. **2022** (2022), <https://doi.org/10.1093/imrn/rnac197>
- [7] M. Baker and C. Petsche, *Global discrepancy and small points on elliptic curves*, Int. Math. Res. Not. **2005** (2005), p. 3791–3834.
- [8] M. Baker and R. Rumely, *Equidistribution of small points, rational dynamics, and potential theory*, Ann. Inst. Fourier **56** (2006), p. 625–688.
- [9] Y. Bilu, *Limit distribution of small points on algebraic tori*, Duke Math. J. **89** (1997), p. 465–476.
- [10] E. Bombieri and U. Zannier, *A note on heights in certain infinite extensions of \mathbb{Q}* , Rend. Mat. Acc. Lincei (9) **12** (2001), p. 5–14.

- [11] G. Burgos, J. Ignacio, P. Philippon, J. Rivera-Letellier and M. Sombra, The distribution of Galois orbits of points of small height in toric varieties, *Amer. J. Math.* **141** (2019), p. 309–381.
- [12] A. Chambert-Loir, Mesures et équidistribution sur les espaces de Berkovich, *J. Reine Angew. Math.* **595** (2006), p. 215–235.
- [13] C. D’andrea, M. Narváez-Clauss and M. Sombra, Quantitative equidistribution of Galois orbits of small points in the N -dimensional torus, *Algebra Number Theory* **11** (2017), p. 1627–1655.
- [14] L. DeMarco and N.M. Mavraki, Variation of canonical height and equidistribution, *Amer. J. Math.* **142** (2020), p. 443–473.
- [15] J. Doyle, P. Fili and B. Tobin, Stochastic Equidistribution and Generalized Adelic Measures, preprint, <https://arxiv.org/abs/2111.08905>
- [16] P. Erdős and P. Turán, On the Distribution of Roots of Polynomials, *Ann. of Math. (2)* **51** (1950), p. 105–119.
- [17] C. Favre, J. Rivera-Letellier, Équidistribution quantitative des points de petite hauteur sur la droite projective, *Math. Ann.* **335** (2006), p. 311–361.
- [18] M. Fekete, Über die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten, *Math. Z.* **17** (1923), p. 228–249.
- [19] W. Gubler, The Bogomolov conjecture for totally degenerate abelian varieties, *Invent. Math.* **169** (2007), p. 377–400.
- [20] M. Langevin, Minorations de la maison et de la mesure de Mahler de certains entiers algébriques, *C. R. Acad. Sci. Paris Sér. I Math.* **303** (1986), p. 523–526
- [21] D. Masser, ”Auxiliary polynomials in number theory”, Cambridge University Press, Cambridge, 2016.
- [22] M. Mignotte, Sur un théorème de M. Langevin, *Acta Arith.* **54** (1989), p. 81–86.
- [23] C. Petsche, A quantitative version of Bilu’s equidistribution theorem, *Int. J. Number Theory* **1** (2005), p. 281–291.
- [24] D. Roy and J. Thunder, An absolute Siegel’s lemma, *J. Reine Angew. Math.* **476** (1996), p. 1–26.

- [25] R. Rumely, On Bilu's equidistribution theorem, *Contemp. Math.* **237** (1999), p. 159–166.
- [26] A. Schinzel, On the product of the conjugates outside the unit circle of an algebraic number. *Acta Arith.* **24** (1973), p. 385–399.
- [27] L. Szpiro, E. Ullmo and S. Zhang, Équirépartition des petits points, *Invent. Math.* **127** (1997), p. 337–347.
- [28] X. Yuan, Big line bundles over arithmetic varieties, *Invent. Math.* **173** (2008), p. 603–649.
- [29] S. Zhang, Small points and adelic metrics, *J. Algebraic Geom.* **4** (1995), p. 281–300.
- [30] S. Zhang, Equidistribution of small points on abelian varieties, *Ann. of Math. (2)* **147** (1998), p. 159–165.