

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Soil fungal communities under slash-and-burn system in Mozambique: A metataxonomic approach

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/2031348> since 2024-11-20T16:07:06Z

Published version:

DOI:10.1111/sum.13062

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Soil fungal communities under slash-and-burn system in Mozambique: A metataxonomic**
2 **approach**

3

4 Dominique Serrani^a, Ilario Ferrocino^b, Cristiana Garofalo^a, Andrea Osimani^a, Maria Rita Corvaglia^b,
5 Vesna Milanović^a, Lucia Aquilanti^a, Valeria Cardelli^{*a}, Andrea Salvucci^a, Stefania Cocco^a, Alves
6 Rafael Rogerio Borguete^c, Giuseppe Corti^{ad}

7

8 ^aDepartment of Agriculture, Food and Environmental Sciences – D3A; Polytechnic University of
9 Marche, 60131 Ancona, Italy.

10 ^bDepartment of Agriculture, Forest and Food Sciences, University of Turin, 10095 Grugliasco, Italy.

11 ^cDepartment of Rural Engineering, Soil Science Division; Faculty of Agronomy and Forestry
12 Engineering, University Eduardo Mondlane. Av. Julius Nyerere, No. 3435, P. Box 257, University
13 Campus, Building #1, Maputo, Mozambique.

14 ^dCREA - Council for Agricultural Research and Analysis of the Agricultural Economy; Centre of
15 Agricultural and Environmental Research, 50125 Firenze, Italy.

16

17

18 *Corresponding author: Department of Agriculture, Food and Environmental Sciences – D3A;
19 Polytechnic University of Marche, 60131 Ancona, Italy. v.cardelli@univpm.it (V. Cardelli).

20

21

22

23

24

25

26

27 **Abstract**

28 This study provides a metataxonomic analysis of the fungal communities in soils under slash-
29 and-burn agroforestry system and offers new insights into the relationships between fungal
30 populations and soil physicochemical features such as pH, the particle size distribution, easily
31 oxidizable organic carbon, total nitrogen, available phosphorus, and the mineralogical composition.
32 Soils from three locations in central Mozambique—Vanduzi, Sussundenga, and Macate—that are
33 subjected to slash and burn were considered to assess the effects of the forest fallow length (temporal
34 variation) and of the land use (charcoal kiln, crop field, and forest; meaning horizontal variation) on
35 the fungal community. The fungi of the genetic horizons (vertical variation) were also considered.
36 Most of the detected fungi were decomposers, antagonists of plant pathogens, and plant-growth
37 promoters; they were differently distributed in relation to the soil's physicochemical properties and
38 the soil use. The variations in the fungi distribution among the locations and between the horizons
39 were considerable, while there were few variations between the different land-use types. The limited
40 differences between land uses indicate the inability of a forest fallow period shorter than 50 years to
41 improve soil fertility and induce changes in the fungal community. The pedological approach used to
42 identify and sample soil horizons allowed us to clearly distinguish the fungal community of the A
43 horizons, those richest in organics and nutrients, and that of the Bo horizons, which have poor fertility.

44

45

46 **Keywords:** soil genetic horizons; Illumina sequencing; land-use change; soil fungi; 26S rRNA gene
47 sequencing

48 **1. Introduction**

49 Slash and burn is a rotational agroforestry system that is widespread in the tropical and
50 subtropical regions of the world (Mertz et al., 2009), where poorly fertile soils such as Oxisols occur
51 (Soil Survey Staff, 2015). In rural areas, farmers use the slash-and-burn approach to a segment of
52 forest to convert it into a cultivated field; to do so, they distribute several charcoal kilns per hectare
53 where stems and branches are used to produce charcoal for the family. The burning activity and the
54 charcoal production produce ashes as a byproduct; these are roughly distributed across the field. Such
55 distribution has the effect of temporarily increasing soil fertility and boosting microbial activity (Gay-
56 des-Combes et al., 2017). After two to four years of cultivation, when crop production is insufficient
57 to maintain family supplies, the field is abandoned and natural reforestation is allowed to occur for
58 decades until the land is cultivated again. Due to the absence of chemical fertilization (Rafael et al.,
59 2018), the forest fallow period is considered essential to restoring a certain level of soil fertility
60 (Gonçalves Lintemani et al., 2019) through soil organic matter (SOM) accumulation and
61 mineralization (Andriamananjara et al., 2020; Silva-Forsberg and Fearnside, 1997). For centuries, the
62 duration of the forest fallow was approximately 50–100 years or more, but the population growth and
63 socio-economic changes that occurred in the second half of the 20th century shortened this period by
64 a significant amount (Chowdhury et al., 2020; Nath et al., 2016), also reducing the ecosystem services
65 provided by forests (Wood et al., 2016). Indeed, studies have established that gradual soil degradation
66 (Gay-des-Combes et al., 2017; Thomaz et al., 2013; Zwartendijk et al., 2020) and the loss of flora,
67 fauna, and microbial diversity (Aguilar-Fernández et al., 2009; Randriamalala et al., 2019) are
68 triggered by the intense deforestation caused by the intensification of the slash-and-burn practice
69 (Curtis et al., 2018).

70 After the effects of a slash-and-burn system on the soil bacterial community in Mozambique were
71 studied (Serrani et al., 2023), it was considered useful to assess the fungal community's diversity in
72 the same context. In fact, as argued by Arévalo-Gardini et al. (2020), due to the influence of microbial

73 activity on ecosystems' stability and fertility, variations in the fungal community may constitute a
74 valid indicator of changes in soil health caused by land management. Fungi are significantly affected
75 by physiographic conditions, environmental contexts related to climate and land management, and
76 soil properties such as the SOM content and fertility level (Shah et al., 2016; Spurgeon et al., 2013;
77 Oehl et al., 2017). Important soil fungi, such as saprotrophic varieties, are fundamentally important
78 decomposers of lignocellulosic remnants (Clocchiatti et al., 2020, van der Wal et al., 2013), while
79 entomopathogenic fungi are endophytes that can enhance plant defenses against harmful insects
80 (Deaver et al., 2019; Vega, 2018). Many studies describe the diversity in the soil fungal population,
81 but few consider fungi in soils submitted to slash and burn. To the best of our knowledge, only
82 Aguilar-Fernández et al. (2009), Adeniyi (2010), Sharmah et al. (2014), and Barraclough and Olsson
83 (2018) have studied fungi variations in soils subjected to slash and burn; these studies have mainly
84 focused on the arbuscular mycorrhizal fungi (AMF) community, a group of fungi that have obligate
85 symbiotic relationships with many plants and which play a specific role in nutrient uptake (e.g.,
86 Deveautour et al., 2018; Yang et al., 2011; Saliou Sarr et al., 2019; Rožek et al., 2020). Improving
87 our knowledge of soil fungal diversity would allow us to understand the complexity of specific
88 ecosystems and their responses to slash-and-burn practice.

89 The aim of this work was therefore to use a metataxonomic approach to evaluate the fungal diversity
90 in the soils of three locations in central Mozambique that are subject to slash and burn, considering
91 the effect of *i*) the three locations as representing different durations of the forest fallow period
92 (temporal variation); *ii*) the land uses forming the slash-and-burn system: charcoal kiln, crop field,
93 and forest (horizontal variation); and *iii*) the development of genetic soil horizons (vertical variation).
94 In so doing, we hypothesized that the soil fungal community can differentiate horizontally and/or
95 vertically, according to land management and soil changes.

96

97 2. Materials and methods

98 2.1. Locations and morphological description of the studied areas

99 Three locations from the Manica province, central Mozambique, were selected: Vanduzi,
100 Sussundenga, and Macate (see Fig. S1 of Supplementary Materials in Serrani et al., 2023). The
101 selected locations all fall into the Agro-Ecological Zone R4, which includes areas between 200 and
102 1000 m above sea level (Maria and Yost, 2006), with the mean annual rainfall ranging from 1000 to
103 1200 mm and the mean annual air temperature around 21°C (Climate Data, 2019). The geology of
104 the zone is dominated by the Mesoproterozoic Southern Irumide Belt (950-1060 Ma), a litho-tectonic
105 unit made up of metamorphic rocks (Chaúque et al., 2019). The predominant soil type belongs to the
106 order of *Oxisols*, which is characterized by primary low fertility and strong erosion due to the
107 topography of the terrain (Maria and Yost, 2006). Furthermore, the soils were recognized as having
108 an aridic moisture regime and a thermic temperature regime (Soil Survey Staff, 2014). Such
109 pedoclimatic conditions have led to the formation of typical tropical woodland (open forest), which
110 is common in the studied locations, comprising savannas and shrublands made up of sparse trees of
111 the leguminous trees *Brachystegia spiciformis* Benth., *Brachystegia tamarindoides* Benth., and
112 *Julbernardia globiflora* (Benth.) with a more or less thick grass understorey of *Themeda triandra*
113 Forssk., *Panicum maximum* Jacq., *Hyparrhenia filipendula* (Hochst.) Stapf, and *Andropogon gayanus*
114 Kunth, referred to as *miombo* (Sitoe, 2004).

115 The three locations were chosen as the slash-and-burn system has been and is still being practised
116 there, but with forest fallow periods of different durations (temporal variation), so as to form the
117 following chronosequence: *i*) in Vanduzi, the forest was ≈ 25 years old, the crop field was 1 year old,
118 and the charcoal kiln was 4 years old; *ii*) in Sussundenga, the forest was ≈ 35 years old, the crop field
119 was 2 years old, and the charcoal kiln was 1 year old; *iii*) in Macate, the forest was ≈ 50 years old, the
120 crop field was 16 years old, and the charcoal kiln was 16 years old (see Table S1 of Supplementary
121 Materials in Serrani et al., 2023). In each location, we took into consideration the soils under the

122 charcoal kiln, the crop field, and forest (horizontal variation), and soil samples from each pedogenic
123 horizon were collected (vertical variation).

124 Further details on the study areas, slash-and-burn systems, and study sites are reported in Serrani et
125 al. (2023).

126

127 2.2. *Sampling campaigns and soil characteristics*

128 After a brief geomorphological and soil survey was conducted in March 2017, the sampling
129 sites were selected in a relatively flat area (plateau) with a gentle slope (2–4%), featuring mostly
130 Oxisols (Soil Survey Staff, 2014) developed from similar metamorphic parent rocks (Chaúque et al.,
131 2019; Wijnhoud, 1997) (see Table S1 of Supplementary Materials in Serrani et al., 2023). To account
132 for the eventual differences in terms of the fungal community across the agricultural seasons, the first
133 sampling campaign was run in March 2017 (autumn) and the second one in November 2017 (spring).
134 For each sampling campaign, soil profiles were opened in a representative area after the preliminary
135 manual opening of mini-pits and auger holes. In the charcoal kilns and agricultural fields, the soil
136 profiles were opened in approximately the middle of their extension, while those in the *miombo* were
137 opened at ≈ 1 m from the trunk of one of the biggest *Brachystegia spiciformis* trees. Once excavated,
138 each profile was described according to Schoeneberger et al. (2012) and sampled according to genetic
139 horizons. In all sites, the studied soils were constituted by a brownish A horizon (umbric) and a
140 reddish Bo (oxic) horizon with a coarse texture, a good degree of aggregation, and the absence of the
141 redoximorphic feature, indicating good drainage and, consequently, a low water-holding capacity
142 (e.g., Agrawal 1991; Suzuki et al., 2007) (see Table S1 of Supplementary Materials in Serrani et al.,
143 2023). About 4 kg of samples were collected from each horizon and stored inside a portable fridge
144 during the field operations. Once in the laboratory, the samples were air-dried and then sieved at 2
145 mm to remove the skeletal particles and coarse vegetal residues.

146 To summarize, for each location (Vanduzi, Sussundenga, and Macate), 12 soil samples were collected
147 (3 land uses x 2 horizons x 2 replicates) from each campaign, for a total of 36 samples.

148

149 2.3. *Soil analyses and microbial DNA extraction and sequencing*

150 The physicochemical and mineralogical analyses run on the soil samples are reported in Table

151 1, which synthesizes information that is fully explained in Serrani et al. (2023).

Table 1. Methods adopted to assess physicochemical properties of each soil sample collected according to locations, land uses, and horizons within Manica province, central Mozambique.

Property	Procedure	Bibliography
pH in water	Potentiometric method, using a combined glass-calomel electrode immersed into the suspension (1:2.5 solid:liquid ratio).	Thomas, 1996
Particle-size distribution	After dissolution of organic cements by Na-hypochlorite (NaOCL)solution at 6% of active chlorine adjusted to pH 9 with HCL, sand (2-0.05 mm) was recovered by wet sieving, while silt (0.05-0.002 mm) was separated from clay (< 0.002 mm) by sedimentation maintaining the columns at 19-20°C.	Lavkulich and Wiens, 1970
Easily oxidizable organic carbon (EOOC)	Walkley-Black method by K-dichromate digestion without application of heating.	Nelson and Sommers, 1996
Total nitrogen (N)	Semi-micro Kjeldahl method.	Bremmer, 1996
Potentially plant-available phosphorous (P)	Olsen method.	Olsen et al., 1954
Mineralogical assemblage	Assessed by X-ray diffractometry on manually compressed powdered samples by using a Philips PW 1830 diffractometer (Fe-filtered Co K α 1 radiation, 35 kV and 25 mA). Minerals were identified on the basis of their characteristic peaks, and a semi-quantitative mineralogical composition was obtained by estimating the area of the diagnostic peaks by multiplying the peak height by its width at half-height.	Brindley and Brown, 1980; Dixon and Schulze, 2002

152

153 Total microbial DNA was extracted from 250 mg of each soil sample using the E.Z.N.A. ® Soil DNA

154 Kit (Omega Bio-Tek, Inc., Georgia, USA), following the manufacturer's instructions. The extracted

155 DNA was quantified using a Qubit dsDNA assay kit (Life Technologies, Milan, Italy) and

156 standardized to 5 ng μL^{-1} . Then, 2.5 μL were used as a template to amplify the D1 domain of the 26S

157 rDNA gene by using the primers and the protocol described by Mota-Gutierrez et al. (2019); a

158 negative control was included in the PCR reactions by replacing the DNA solution with water. The

159 26S gene region provides a higher alpha diversity index and greater fungal rRNA taxonomic depth

160 and robustness results compared with ITS2 (Mota-Gutierrez et al., 2019). The PCR amplicons were

161 purified, tagged, and sequenced according to the Illumina metagenomic pipeline instructions. The
162 sequencing was performed using a MiSeq Illumina instrument (Illumina, San Diego, USA) with V3
163 chemistry and generated 2x250 bp paired-end reads, according to the manufacturer's instructions.

164

165 *2.4. Bioinformatic analysis*

166 After sequencing, reads were analyzed using the Quantitative Insights into Microbial Ecology
167 QIIME2 (Bolyen et al., 2019). Primers and adapters were trimmed using Cutadapter and then filtered
168 for quality using the DADA2 algorithm (Callahan et al., 2016), removing low-quality bases and
169 chimeric sequences with the DADA2 denoise-paired plug-in of QIIME2. A total of 3.820.038 clean
170 reads were used for downstream analysis (99% of the sample coverage). Amplicon Sequence Variants
171 (ASVs) generated by DADA2 were used for a taxonomic assignment using the QIIME feature-
172 classifier plug-in against the SILVA-implemented database for fungi (Mota-Gutierrez et al., 2019).
173 Briefly, the database was obtained using the large subunit rRNA gene sequences from the Silva
174 database and from NCBI. The fungi taxonomic assignment was double-checked using BLAST suite
175 tools. The QIIME2 diversity script was used to perform alpha diversity analysis. The data generated
176 by sequencing were deposited in the NCBI Sequence Read Archive (SRA) and are available under
177 the Bioprojects Accession Number PRJNA631872: biosample accession numbers from
178 SAMN14895437 to SAMN14895491 and from SAMN14895517 to SAMN14895548.

179

180 *2.5. Statistical treatment of the data*

181 Statistical analyses of the soil's physicochemical properties are reported in Serrani et al. (2023)
182 and briefly reported below. Physicochemical soil data were statistically treated using the R program
183 (vv 1.3.1093) workspace. ANOVA was used to test the similarity of the two sampling campaigns for
184 physicochemical soil properties [pH, particle size distribution, easily oxidizable organic carbon
185 (EOOC), total N, and available P] (see Table S2 of Supplementary Materials in Serrani et al., 2023;
186 $P > 0.05$). Once it was confirmed that the samples collected in the two sampling campaigns were

187 replicates, ANOVA was run to test significative differences for sampling locations, land uses, and
188 horizons (see Table S3 of Supplementary Materials in Serrani et al., 2023; $P > 0.05$). The contrasted
189 results of the whole profiles were obtained from the weighted mean of each outcome for the thickness
190 of the A and Bo horizons of each soil sample. To apply the parametric test, we verified the normal
191 distribution of the data using the Shapiro–Wilk statistical test (stats R package) and the equal
192 variances using Levene's test (car R package), both at a 5% of significance level. When the data were
193 non-normally distributed, each numerical variable was transformed using the Box–Cox procedure
194 (Meloun et al., 2005). When the normality assumption was validated, a post hoc Tukey's Honest
195 Significant Difference (HSD) test with $P \leq 0.05$ was used to compare the means; in contrast, the
196 Kruskal–Wallis non-parametric test was used to assess the significance of the differences. In the case
197 of heteroscedasticity, the Welch one-way ANOVA test was used ($P \leq 0.05$), while, in case of
198 heteroscedasticity and non-normality, the Friedman test (rstatix package) combined with Kendall's
199 W were used to measure the Friedman test effect size and pairwise Wilcoxon signed-rank tests.
200 Fungal α -diversity was assessed using the Chao1 index and the Shannon diversity index, calculated
201 using the diversity function of QIIME2 on an ASV table rarefied at the lowest feature count/sample.
202 A Bray–Curtis distance matrix was generated by QIIME2 and used to build the principal coordinate
203 analysis (PCoA) and to perform PERMANOVA as a function of location, land use, or horizon.
204 Variables that were not normally distributed were presented as the median (interquartile range).
205 Variables were compared using the Mann–Whitney U test or the Kruskal–Wallis test, as appropriate.
206 An ASVs table was then imported in R to build the heatmap using the *made4* function. Spearman
207 correlation analysis between physicochemical properties and fungi was performed with
208 the *psych* package and *corrplot()* from *corrplot* R package. The P values were adjusted for multiple
209 testing using the Benjamini–Hochberg procedure, which assesses the false discovery rate (FDR).
210 The arithmetic means and relative standard deviations for physicochemical properties (see Tables S4,
211 S5, and S6 of the Supplementary Materials in Serrani et al., 2023) and ASVs were calculated for the

212 sampling locations (n=12), total land use (n=12), land use of each area (n=4), total horizons (n=18),
213 and the horizon of each site (n=6).

214

215 **3. Results and discussion**

216 *3.1. Fungal diversity*

217 Differences in fungal composition as a function of location, land use, and horizons were
218 examined at the highest taxonomic resolution reached, namely, at the genus or family level (Fig. 1).
219 ASVs were detected in the dataset and grouped into two main clusters: *i*) Cluster 1 was characterized
220 by the highest frequency of *Sarcinomyces*, *Catenulifera*, *Chaetomium*, *Zygoascus*, *Fusarium*,
221 *Trichoderma*, and *Chaetomiaceae* and included most of the samples from the Macate and Vanduzi
222 soils and the A horizons; *ii*) Cluster 2 exhibited the highest frequency of *Aureobasidium*,
223 *Cladosporium*, *Malassezia*, *Pichia*, *Aspergillus*, *Saccharomyces*, and *Acremonium* and included most
224 of the samples from the Sussundenga soils and the Bo horizons. Toju et al. (2016) analyzed the fungal
225 network in a cool–temperate forest in Hokkaido (Japan) and found that, as in our case, *Malassezia*
226 and *Cladosporium* had a strong preference for the B horizons. Elsewhere, Chen et al. (2019) found
227 an abundance of saprotroph fungi in organic soil and an abundance of symbiont fungi in the mineral
228 topsoil under a subtropical forest.

229 Analysing the alpha diversity values as a function of the locations, we observed the highest levels of
230 richness (the Shannon and Chao1 indices) and ASVs in the Macate soils (FDR<0.05, data not shown).
231 Alpha diversity as a function of land use did not show significant differences, while the comparison
232 between horizons highlighted greater complexity in the A horizon than in the Bo horizon (FDR<0.05,
233 data not shown). The alpha diversity comparison referring to horizons showed the fungi to decrease
234 in both number and species from the A to the Bo horizons, that is, with increasing depth; this finding
235 was also reported by Warcup (1951) and Jumpponen et al. (2010). Some studies have reported that
236 soils with high fungal richness and diversity showed a relatively large content of N (Mueller et al.,

237 2014; Weber et al., 2013), and this could explain why the highest Shannon and Chao1 indexes were
238 found in the Macate soils and in the A horizons, where relatively high total N content occurred (see
239 Tables S4 and S6 of the Supplementary Materials in Serrani et al., 2023). Since the physicochemical
240 changes occurring along the soil profile induce the development of spatial niches that are able to
241 accommodate different fungal communities (Chen et al., 2019), we assume this happened in our soils
242 where, from the A to the Bo horizons, nutrients decreased and roots increased (see Tables S1 and S6
243 of the Supplementary Materials in Serrani et al., 2023).

244

245 3.2. *Effect of location (temporal variation) on fungal diversity*

246 PCoA based on the Bray–Curtis distance matrix showed a partial overlapping of fungi for the
247 Vanduzi and Sussundenga soils, which were separated from the Macate soils (Fig. S1 of
248 Supplementary Materials, $P < 0.001$). Considering the relative frequency across locations, Vanduzi
249 soils showed the highest frequency of *Cladosporium* (Fig. 2, $FDR < 0.05$), while Macate soils were
250 characterized by the highest frequencies of *Catenulifera*, *Fusarium*, *Penicillium*, *Sarcinomyces*,
251 *Trichoderma*, and *Zygoascus* (Fig. 2, $FDR < 0.05$).

252 *Cladosporium* is a genus that includes 993 heterogeneous and ubiquitous kinds of hyphomycetes that
253 are well-known as common endophytes (Bensch et al., 2012). Several bioactive molecules that are
254 active against bacteria and fungi have been isolated from endophytic *Cladosporium* species, thus
255 indicating the main role of this group of fungi in producing antimicrobial compounds that are involved
256 in the control of plant pathogens (Yehia et al., 2020). Additionally, because of the generally higher N
257 content, the highest prevalence of *Cladosporium* is often associated with intensive cultivation
258 systems, but it has been also reported to be an important taxon of the phyllosphere microbial
259 community (Abdelfattah et al., 2016). *Cladosporium* is also involved in plant P absorption (Shi et al.,
260 2020), and the highest frequency of these fungi were likely present in the Vanduzi soils due to the
261 highest content levels of available P in these soils (see Table S4 of Supplementary Materials in Serrani

262 et al., 2023), despite no significant correlation to the soil's physicochemical properties being found
263 (Fig. 3).

264 The genus *Catenulifera* includes anamorph species of *Hyphodiscus*, a genus of discomycetes that has
265 been found to be associated with decaying wood and the fruit bodies of other fungi (Bogale et al.,
266 2010; Hosoya et al., 2002). As far as we know, no information exists about the interaction between
267 *Catenulifera* requirements and soil properties; however, based on the correlation plot (Fig. 3),
268 *Catenulifera* appears to be negatively related to pH and sand, which displayed the lowest values in
269 the Macate soils. This led us to hypothesize that *Catenulifera* were abundant in Macate because they
270 prefer soil environments with acidophilic reactions and relatively high contents of silt and clay;
271 however, the Macate soils also showed the highest levels of organics and N content (see Table S4 of
272 Supplementary Materials in Serrani et al., 2023), even though no significant correlation between
273 *Catenulifera* and these parameters was found (Fig. 3).

274 *Fusarium* is a genus of saprotrophic fungi and/or fast-growing colonizers of the rhizosphere in
275 response to plant exudates (Goncharov et al., 2020), and its proliferation through the formation of
276 both macroconidia and ascospores may be favored by soil moisture conditions (Lemmens et al.,
277 2004). In our soils, the positive correlation of *Fusarium* with total N and EOO (Fig. 3), which
278 abounded in the Macate soils, was ascribed to the accumulation of decaying organic matter provided
279 by the mulching in the crop field and by the presence of relatively well-developed and poorly
280 disturbed litter in the forest, with both mulching and dense forest being able to maintain a certain
281 level of soil moisture.

282 *Penicillium* is a common soil fungi genus that includes plant-beneficial microorganisms (Altaf et al.,
283 2018; Das et al., 2021, Efthymiou et al., 2018a,b); it is also known for growing in extreme
284 environments, including highly acidic soils (Diao et al., 2019; Yadav et al., 2019; Warcup, 1951).
285 These properties of *Penicillium* effectively explain our results, since the ASVs of this genus showed
286 an inverse relation with pH (Fig. 3), which was the lowest in the Macate soils.

287 As *Sarcinomyces* endophytic fungi, their highest frequency in the soils of Macate was ascribed to the
288 relatively pronounced presence of decaying organic matter due to mulching (in the crop field) and
289 forest development, as also reported by Li et al. (2018). Moreover, the correlation plot showed that
290 the frequency of *Sarcinomyces* was inversely correlated to available P and sand (Fig. 3). Given that
291 there are no specific indications for *Sarcinomyces*, since endophytic fungi are often isolated from
292 sandy soils as they produce growth-promoting metabolites that help the host plants to survive under
293 soil stress conditions, the observed inverse correlation was ascribed to these fungi's general resistance
294 to drought and salinity (e.g., Hamayun et al., 2010; Khan et al., 2012, 2016).

295 Species belonging to the genus *Trichoderma* are considered plant-growth promoters, biocontrol
296 agents (Ji et al., 2020; Oskiera et al., 2017; Zhang et al., 2020a), and improvers of N and P availability
297 as they increase the activity of urease, phosphatase, catalase, and cellulase (Ji et al., 2020; Makhuvele
298 et al., 2017). This evidence is aligned with the positive correlation of these fungi with EEOC and
299 total N (Fig. 3), which were abundant in the soils of Macate.

300 Members of the genus *Zygoascus* have been reported to play a role as biofertilizers since they can
301 solubilize soil phosphates (Das et al., 2021). The highest abundance of *Zygoascus* in the Macate soils
302 and their positive correlation with EEOC, total N, and available P (Fig. 3) allowed us to hypothesize
303 that they have a preference for soil niches enriched with organic matter.

304 To summarize, the partial overlapping of fungi in the Vanduzi and Sussundenga soils suggested the
305 irrelevance of the different durations of the forest fallow period for these two locations: 25 and 35
306 years, respectively. In the Macate soils, the different fungal compositions and the highest ASV
307 abundance were probably favored by the higher levels of nutrients, which are due to the pedogenic
308 conditions and soil management (mulching in the crop field, few disturbances in the forest), rather
309 than the different forest fallow period. Considering the fungi's ecosystem/ecological functions (Table
310 2), the most abundant fungi in the Macate soils played important roles, acting as decomposers,
311 antagonists of plant pathogens, and plant-growth promoters.

Table 2. The most abundant fungi (ASVs) and their ecosystem/ecological functions in soils according to locations, land uses, and horizons within Manica province, central Mozambique. Abundances significantly differ at $FDR \leq 0.05$.

	Clade	Family/Genus	Ecosystem/ecological functions	Bibliography
Location				
Vanduzi	Ascomycota	<i>Cladosporium</i>	Antagonistics of plant pathogens; litter and wood saprotrophs; foliar endophytes	Frac et al., 2018; Yehia et al., 2020
Macate	Ascomycota	<i>Catenulifera</i>	Decomposers and fungi pathogens; wood saprotrophs	Bogale et al., 2010; Frac et al., 2018; Hosoya et al., 2002
		<i>Fusarium</i>	Soil-borne root pathogenic fungi; litter saprotrophs; foliar endophytes	Frac et al., 2018; Ge et al., 2021a,b; Goncharov et al., 2020
		<i>Penicillium</i>	Increase in fertilized soils; antagonistics of plant pathogens; plant-growth promoters; litter saprotrophs; foliar endophytes	Altaf, 2018; Das et al., 2021; Efthymiou et al., 2018a,b; Frac et al., 2018
		<i>Sarcinomyces</i>	Ectomycorrhizal mutualists; adapted to high temperatures and low water activity; litter saprotrophs; foliar endophytes	Li et al., 2018; Sterflinger, 1998; Volkmann et al., 2003
		<i>Trichoderma</i>	Increase in fertilized soils; antagonistics of plant pathogens; improvers of the plant health and root growth; litter saprotrophs; foliar endophytes	e.g. Frac et al., 2018; Oskiera et al., 2017; Vinale et al., 2008
	Saccharomyceta	<i>Zygoascus</i>	Biofertilizers	Das et al., 2021; Frac et al., 2018
Land use				
Crop field	Sordariales	<i>Chaetomium</i>	On cellulose-rich soil materials or on dung; adapted to arid climate; litter saprotrophs; foliar endophytes	Ahmed et al., 2016; Frac et al., 2018
Forest	Sordariales	<i>Chaetomium</i>	On cellulose-rich soil materials or on dung; adapted to arid climate; litter saprotrophs; foliar endophytes	Ahmed et al., 2016; Frac et al., 2018
	Ascomycota	<i>Penicillium</i>	Increase in fertilized soils; antagonistics of plant pathogens; plant-growth promoters; litter saprotrophs; foliar endophytes	Altaf, 2018; Das et al., 2021; Efthymiou et al., 2018a,b; Frac et al., 2018
Horizon				
A	Dothideomycetes	Aureobasidiaceae	Fungal endophytes. Dominant in forest soil; decomposers in agricultural soils; increase after nitrogen fertilization	Frac et al., 2018; Khan et al., 2016
	Sordariales	Chaetomiaceae	Antagonistics of plant pathogens; degraders of complex SOM	Chovanova and Zamocky, 2016; Frac et al., 2018; Mohammed et al., 2019
		<i>Chaetomium</i>	On cellulose-rich soil materials or on dung; adapted to arid climate; litter saprotrophs; foliar endophytes	Ahmed et al., 2016; Frac et al., 2018
	Saccharomyceta	<i>Meyerozyma</i>	Involved in the solubilization of phosphates and xylose fermentation; antagonistics of plant pathogens; epiphytes; foliar endophytes	Arumugam et al., 2020; Frac et al., 2018; Kim et al., 2016; Nakayan et al., 2013; Procópio and Barreto, 2021
	Tremellomycetes	Mrakiaceae	Adapted to low temperatures; nitrate and nitrite utilizers; dominant in forest soil; saprotrophic and parasitic fungi	Frac et al., 2018; Sannino et al., 2020; Zhang et al., 2020b
<i>Papiliotrema</i>		Contains species able to interact with AMF† to improve plant N and P uptake; litter saprotrophs; mycoparasites; fungal decomposers	Frac et al., 2018; Leguina et al., 2019	

		<i>Trichoderma</i>	Increase in fertilized soils; antagonistics of plant pathogens; improvers of the plant health and root growth; litter saprotrophs; foliar endophytes	e.g. Frac et al., 2018; Oskiera et al., 2017; Vinale et al., 2008
	Saccharomyceta	<i>Zygoascus</i>	Biofertilizers	Das et al., 2021; Frac et al., 2018
Bo	Saccharomyceta	Debaryomycetaceae	Involved in xylose fermentation; able to produce bioethanol	Arumugam et al., 2020; Hui et al., 2014
	Basidiomycota	<i>Malassezia</i>	Able to colonize a wide range of habitats, including oligotrophic soils; soil saprotrophs; root-associated fungi	Amend, 2014; Frac et al., 2018; Toju et al., 2016
	Sordariomycetes	Microascaceae	Saprobic and plant pathogens; decomposers in agricultural soils; increase after nitrogen fertilization	Frac et al., 2018; Sandoval-Denis et al., 2016
	Saccharomyceta	<i>Pichia</i>	Phosphate- and zinc-solubilizers; thermotolerant yeasts; involved in xylose fermentation; able to produce bioethanol; antagonistics of plant pathogens	Frac et al., 2018; Chamnipa et al., 2018; Kumla et al., 2020; Pongcharoen et al., 2018; Procópio and Barreto, 2021
		<i>Saccharomyces</i>	Phosphate- and zinc-solubilizers; thermotolerant yeasts; involved in glucose fermentation; able to produce bioethanol	Frac et al., 2018; Kumla et al., 2020; Pongcharoen et al., 2018;

†AMF = Arbuscular micorrhizal fungi

312

313 3.3. Effect of land-use (horizontal variation) on fungal diversity

314 The PCoA showed a partial overlapping of fungi as function of land use (Fig. S2, $P < 0.001$).

315 Comparing the different land uses, only 2 out of 37 fungi showed different ASV distributions,

316 *Chaetomium* and *Penicillium*, which showed the highest frequencies in both crop fields and forest

317 soils (Fig. 4, $FDR < 0.05$). Since the samples were small, the frequency of *Penicillium* for crop field

318 soils was slightly higher than that of the charcoal kiln soils, but this was not statistically different.

319 The *Chaetomium* genus belongs to the *Chaetomiaceae* family and is known to be a producer of

320 antimicrobial metabolites against plant pathogens, including fungi and insects (Chovanova and

321 Zamocky, 2016; Mohammed et al., 2019). The *Chaetomiaceae* family is also linked to the

322 degradation of complex SOM (Paula et al., 2020); in particular, the genus *Chaetomium* abounds in

323 soils rich in cellulosic biomass because of the cellulose-degrading capabilities of this genus's

324 members (Ahmed et al., 2016). These characteristics effectively explain why the highest frequency

325 of *Chaetomium* was found in crop fields and forest soils, where they are likely favored by the presence

326 of crop residues and litter accumulation (Ahmed et al., 2016, Soyong et al., 2001). The highest

327 *Penicillium* distribution in the soils under forests (and crop fields) can be ascribed to their adaptability

328 to the low pH values that characterized these soils, as previously reported and as suggested by the
329 correlation analysis (Fig. 3).

330 *Papiliotrema* was the only taxon associated with the charcoal kiln soils of Macate (Fig. S3 of
331 Supplementary Materials, FDR<0.05). Few data are available for *Papiliotrema*, formerly
332 *Cryptococcus*. Members of this genus were found to be predominant in rice-storage granaries (Shi et
333 al., 2021) while, in soil, *Papiliotrema laurentii* was observed to develop a synergic interaction with
334 AMF to improve plants' uptake of N and P (Leguina et al., 2019) and the solubilization of scarcely
335 soluble forms of phosphate (apatites) and zinc (ZnO and ZnCO₃) (Kumla et al., 2020). The presence
336 of *Papiliotrema* in the charcoal kiln soils of Macate can be attributed to the possible presence of
337 phosphatic minerals, which could have been generated by repeated combustions in the same area; the
338 higher availability of P in this soil supports this hypothesis.

339 However, the low number of variations in the fungal composition between charcoal kilns, crop fields,
340 and forests indicate that the different land uses had little influence.

341

342 3.4. Effect of the horizon (vertical variation) on fungal diversity

343 As reported above, fungi tend to create distinct networks throughout the soil; indeed, a certain
344 degree of separation of fungi between the A and Bo horizons was highlighted by the PcoA (Fig. S4,
345 $P<0.001$). The ASVs that were mainly associated with the A horizons were *Aureobasidiaceae*,
346 *Chaetomiaceae*, *Chaetomium*, *Meyerozyma*, *Mrakiaceae*, *Papiliotrema*, *Trichoderma*, and *Zygoascus*
347 (Fig. 5, FDR<0.05), while the Bo horizons displayed the strongest association with
348 *Debaryomycetaceae*, *Malassezia*, *Microascaceae*, *Pichia*, and *Saccharomyces* (Fig. 5, FDR<0.05).

349 For the A horizons, members of *Aureobasidiaceae*, *Chaetomiaceae*, and *Chaetomium* are endophytic
350 fungi that are particularly abundant in leaves and stems (Habtewold et al., 2020, Khan et al., 2016),
351 whereas *Meyerozyma*, *Trichoderma*, and *Zygoascus* were found to play a role in the soil in the
352 solubilization of phosphates (Gizaw et al., 2017; Kim et al., 2016; Saravanakumar et al., 2013). These
353 reports agree somewhat with the correlations we found, which highlighted the following positive

354 relations: *i) Aureobasidiaceae* and *Trichoderma* with EEOC and total N, *ii) Chaetomiaceae* with
355 available P, and *iii) Zygoascus* with EEOC, total N, and available P (Fig. 3). The fungi belonging to
356 the *Mrakiaceae* family are known to be able to adapt their physiology to low temperatures (Sannino
357 et al., 2020) because of their ability to use nitrates and nitrites and to produce enzymes such as lipases,
358 amylases, proteases, pectinases, cellulases, and chitinases, and ligninolytic enzymes (Zhang et al.,
359 2020b). The presence of *Mrakiaceae* in the A horizons of all the soils indicated that not all of the
360 members of this family are adapted to cold environments and that their ability to produce a broad
361 spectrum of degradative enzymes enables these fungi to perform well where organic matter abounds.
362 Therefore, it appeared that the group of fungi associated with the A horizons was favored by the
363 abundance of organic matter and nutrients and, from an ecosystem/ecological point of view (Table
364 2), that they play the roles of decomposers, antagonists of plant pathogens, and plant-growth
365 promoters.

366 For the Bo horizons, the associated fungi showed inverse relations for *Debaryomycetaceae* and
367 *Saccharomyces* with EEOC and total N, *Malassezia* with total N, and *Pichia* with available P.
368 Although little information is available for *Debaryomycetaceae*, this family of yeasts is involved in
369 the xylose fermentation of biomass with the potential to produce bioethanol (Hui et al., 2014;
370 Arumugam et al., 2020). They probably abound in the Bo horizons due to their large number of roots,
371 which constitute a lignocellulosic substrate from which xylose can be freed during root decay (e.g.,
372 Cheshire et al., 1990; Machinet et al., 2009) and which excrete exudates containing xylose (e.g.,
373 Graystone and Campbell, 1996). For *Saccharomyces*, the wild species are commonly associated with
374 tree substrates (bark, leaves, exudates, and litter) and soil (Alsammar and Delneri, 2020), but they are
375 also known as *i) siderophore* producers in both bulk and rhizosphere soils, and *ii) being* responsible
376 for various processes that are beneficial to plants (Das et al., 2021). Because of this, the presence of
377 *Saccharomyces* in the Bo horizons was ascribed to the conspicuous quantity of roots in the sub-
378 surface horizons of several soils (those under the charcoal kilns of Sussundenga and Macate and the
379 forests of Vanduzi and Macate). *Malassezia* can colonize a wide range of extreme habitats (Amend,

2014), but Toju et al. (2016) found that *Malassezia* diffused in soil, especially in the Bo horizons, as in our case. The inverse relationship with total N and the abundance in the Bo horizons indicate that these fungi prefer soil environments with low fertility. Some fungi belonging to the *Pichia* genus are known to be able to produce siderophores and, similarly to *Saccharomyces*, to solubilize zinc and phosphates (Kumla et al., 2020; Nakayan et al., 2009), thus explaining the inverse relation with available P. In addition, the *Pichia* genus was recognized for its ability to ferment xylose and produce bioethanol (Arumugam et al., 2020, Chamnipa et al., 2018; Pongcharoen et al., 2018), as was the case for *Debaryomycetaceae*; because of this, their larger concentrations in the Bo horizons were explained by the notable presence of roots. *Microascaceae* is a scarcely known fungi taxon that includes saprobes and plant pathogens (Sandoval-Denis et al., 2016) and degraders of labile organics (Lueders et al., 2006; Zhang et al., 2018). This taxon showed no correlation with the analytical parameters, but its abundance in the Bo horizons was explained by the large number of roots generally present in these horizons, which provided suitable organic materials during root decay. Therefore, the group of fungi harboring the Bo horizons is probably favored by the presence of many roots and the oligotrophic conditions; it also appeared to be involved in xylose fermentation (Table 2).

No significant difference was observed for the vertical variation among land uses, while several differences were observed between the fungi and soil horizons within each location. In Vanduzi, *Chaetomiaceae*, *Meyerozima*, *Papiliotrema*, and *Zygoascus* were associated with the A horizons, while *Debaryomycetaceae*, *Malassezia*, and *Pichia* were associated with the Bo horizons (Fig. S5 of Supplementary Materials, FDR<0.05). At Sussundenga, *Trichoderma* and *Zygoascus* prospered in the A horizons (Fig. S6 of Supplementary Materials, FDR<0.05); meanwhile, in Macate, *Aureobasidiaceae* and *Chaetomiaceae* were the most abundant in the A horizons, with *Catenulifera*, *Malassezia*, and *Microascaceae* predominating in the Bo horizons (Fig. S7 of Supplementary Materials, FDR<0.05). As mentioned before, *Chaetomiaceae*, *Aureobasidiaceae*, *Meyerozima*, *Trichoderma*, and *Zygoascus* were related to the presence of SOM and available nutrients, and their abundance in the A horizons is explained by their relatively high contents of EEOC, total N, and

406 available P. On the contrary, *Debaryomycetaceae*, *Malassezia*, and *Pichia* seemed to prefer
407 oligotrophic environments, which, in these soils, are represented by the Bo horizons. The tendency
408 of *Microascaceae* to proliferate in presence of roots may be explained by the significant amount of
409 living and dead roots observed in the Bo horizons in Macate (see Table S1 of the Supplementary
410 Materials in Serrani et al., 2023). In summary, the fungal community exhibited significant vertical
411 variations with a clear separation between the A and Bo horizons, which was mostly driven by the
412 distribution of nutrients and roots.

413

414 3.5. Correlation between fungi and the soil's physicochemical properties

415 In addition to the above-mentioned correlations, Fig. 3 shows the other relationships between fungi
416 and soil properties in the studied soils. Specifically, inverse relations were observed for *Acremonium*
417 and pH, and for *Plectosphaerellaceae* and *Ogataea* with clay content (FDR<0.05), whereas
418 *Aureobasidium* showed a negative relation with silt and a positive relation with clay content
419 (FDR<0.05).

420 Members of the genus *Acremonium* include plant pathogens, wood saprotrophs, and mycoparasitic
421 species (Nguyen et al., 2016), and they were found to be more abundant in N- and P-fertilized soils
422 with an acidic pH (4.6 and 4.8) than in soils with a higher pH (Zhou et al., 2016). These findings
423 agree with the results of our corrplot (Fig. 3).

424 The *Plectosphaerellaceae* family comprises numerous plant pathogen genera and soil-borne species
425 that have been detected in sandy and loamy soils (Giraldo and Crous, 2019), thus explaining the
426 inverse correlation between *Plectosphaerellaceae* and clay. The genus *Ogataea* is characterized by
427 thermotolerant and nitrate-assimilating methylotrophic yeasts (Limtong et al., 2008; Suh and Zhou,
428 2010) that are probably more suited to coarse-textured soils where nitrate availability is higher than
429 in clay-rich soils.

430 *Aureobasidium* is a genus of hyphomycetes fungi that inhabit various extreme environments (Bozoudi
431 et al., 2018; Zalar et al., 2008), including the stones and rocks of moderate or humid climates

432 (Sterflinger, 2010). This suggests that their distribution in our soils was enabled by the fine separates,
433 possibly because of their ability to retain humidity.

434

435 **4. Conclusions**

436 This study provides one of the first metataxonomic analyses of soil-associated fungi in soils
437 undergoing slash-and-burn practices and offers new insights into the relationship between fungal
438 populations and soil physicochemical properties. The results highlighted the separation of fungi into
439 two main groups: those affected by temporal, spatial, and vertical soil variations and those that are
440 homogeneously distributed in all the investigated soils. Within these diverse abundances, the main
441 differences were found among locations and between horizons. In the first case, the fungal
442 distribution was ascribed to genetic soil properties and soil management rather than to the different
443 lengths of the forest fallow period; in the second case, the ecological pressures responsible for fungal
444 differentiation were recognized in the different dotation of SOM, nutrients, and living and decaying
445 roots between the A and Bo horizons. In contrast, land use exerted negligible influence in determining
446 differences in the fungal community, especially for the soils under crop fields and forests. Our
447 findings indicate that temporal, horizontal, and vertical fungal distributions mainly depend on soil
448 genesis and management, and that forest fallow is ineffective in producing substantial changes in the
449 fungal community and, consequently, recovering soil biological fertility. Because of this, the fungi
450 harbored in different soil environments have the potential to be considered ecosystem/ecological
451 indicators of soil conditions and health. However, even though our approach is a commonly used
452 method for characterizing fungi, we are aware of the limitations of our study. For example, the
453 extraction method is not equally efficient for the different forms of fungi (simple cell, hyphae, or
454 spore) and significant variation exists between methods. Thus, the metataxonomic technique may
455 lead to possible biases due to amplification and may produce over- or underestimations of ASVs. In
456 addition, sexual or asexual forms can produce different classifications in taxonomy. Because of this,

457 we believe that additional studies are needed to further disclose the role of fungi in various soil
458 horizons and the role of well-differentiated soil horizons in stimulating the proliferation of useful
459 fungi.

460

461 **Acknowledgments**

462 The authors are indebted to the Sussundenga Research Station (*Instituto de Investigação Agrária de*
463 *Moçambique*). Grants were received through the Universidade Eduardo Mondlane by the Italian
464 Cooperation and Development Agency for the project “Applied Research and Multi-sectorial
465 Program” (No. 5.2.1), and by the project “PSA2017-Discovering “terra preta” in Mozambique: a
466 model for sustainable agroforestry systems to preserve soil, forest, and wilderness areas”, which was
467 funded by the Polytechnic University of Marche (Italy). The findings, conclusions, and
468 recommendations presented in this document do not necessarily reflect the views of the donors.

469

470 **References**

- 471 Abdelfattah, A., Wisniewski, M., Li Destri Nicosia, M. G., Cacciola, S. O., & Schena, L. (2016).
472 Metagenomic analysis of fungal diversity on strawberry plants and the effect of management
473 practices on the fungal community structure of aerial organs. *PLoS ONE*, *11*, e0160470.
474 <https://doi:10.1371/journal.pone.0160470>
- 475 Adeniyi, A. S. (2010). Effects of slash and burning on soil microbial diversity and abundance in the
476 tropical rainforest ecosystem, Ondo State, Nigeria. *African Journal of Plant Science*, *4*, 322-329.
- 477 Agrawal, R. P. (1991). Water and nutrient management in sandy soils by compaction. *Soil & Tillage*
478 *Research*, *19*(2), 121-130. [https://doi.org/10.1016/0167-1987\(91\)90081-8](https://doi.org/10.1016/0167-1987(91)90081-8)
- 479 Aguilar-Fernández, M., Jaramillo, V. J., Varela-Fregoso, L., & Gavito, M. E. (2009). Short-term
480 consequences of slash-and-burn practices on the arbuscular mycorrhizal fungi of a tropical dry
481 forest. *Mycorrhiza*, *19*, 179–186. <https://doi.org/10.1007/s00572-009-0229-2>

482 Ahmed, S. A., Khan, Z., Wang, X. -W., Moussa, T. A., Al-Zahrani, H. S., Almaghrabi, O. A., Sutton,
483 D. A., Ahmad, S., Groenewald, J. Z., Alastruey-Izquierdo, A., van Diepeningen, A., Menken, S.
484 B. J., Najafzadeh, M. J., Crous, P. W., Cornely, O., Hamprecht, A., Vehreschild, M. J. G. T.,
485 Kindo, A. J., & de Hoog, G. S. (2016). Chaetomium-like fungi causing opportunistic infections
486 in humans: a possible role for extremotolerance. *Fungal Diversity*, 76, 11-26.
487 <https://doi.org/10.1007/s13225-015-0338-5>

488 Alsammar, H., & Delneri, D. (2020). An update on the diversity, ecology and biogeography of the
489 *Saccharomyces* genus. *FEMS Yeast Research*, 20, 1-12. <https://doi.org/10.1093/femsyr/foaa013>

490 Altaf, M. M., Imran, M., Abulreesh, H. H., Khan, M. S. S., & Ahmad, I. (2018). Diversity and
491 applications of *Penicillium* in plant-growth promotion. In V. K. Gupta, & S. Rodriguez-Couto
492 (Eds.), *New and Future Developments in Microbial Biotechnology and Bioengineering:*
493 *Penicillium system properties and applications* (pp. 261-276). Elsevier.
494 <http://dx.doi.org/10.1016/B978-0-444-63501-3.00015-6>

495 Amend, A. (2014). From Dandruff to Deep-Sea Vents: *Malassezia*-like fungi are ecologically hyper-
496 diverse. *PloS Pathogens*, 10. <https://doi.org/10.1371/journal.ppat.1004277>

497 Andriamananjara, A., Hewson, J., Razakamanarivoa, H., Andrisoa, R. H., Ranaivosona, N.,
498 Ramboatiana, N., Razafindrakoto, M., Ramifehiarivo, N., Razafimanantsoa, M. -P., Rabeharisoa,
499 L., Ramananantoandro, T., Rasolohery, A., Rabetokotany, N., & Razafimbelo, T. (2020). Land
500 cover impacts on aboveground and soil carbon stocks in Malagasy rainforest. *Agriculture,*
501 *Ecosystem & Environment*, 233, 1-15. <http://dx.doi.org/10.1016/j.agee.2016.08.030>

502 Arévalo-Gardini, E., Canto, M., Alegre, J., Arévalo-Hernández, C. O., Loli, O., Julca, A., Baligar, V.
503 (2020). Cacao agroforestry management systems effects on soil fungi diversity in the Peruvian
504 Amazon. *Ecological Indicators*, 115, 106404. <https://doi.org/10.1016/j.ecolind.2020.106404>

505 Arumugam, N., Boobalan, T., Saravanan, S., Jothi Basu, M., Arun, A., Suganya Devi, T., Kavitha,
506 T. (2020). In *silico* and in *vitro* comparison of nicotinamide adenine dinucleotide phosphate

507 dependent xylose reductase rossmaan fold in *Debaryomycetaceae* yeast family. *Biocatalysis and*
508 *Agricultural Biotechnology*, 24, 101508. <https://doi.org/10.1016/j.bcab.2020.101508>

509 Barraclough, A. D., Olsson, P. A. (2018). Slash-and-burn practices decrease arbuscular mycorrhizal
510 fungi abundance in soil and the roots of *Didierea madagascariensis* in the dry tropical forest of
511 Madagascar. *Fire*, 1(3), 37. <https://doi.org/10.3390/fire1030037>

512 Bensch, K., Braun, U., Groenewald, J. Z., Crous, P. W. (2012). The genus *Cladosporium*. *Studies in*
513 *Mycology*, 72, 1-401. <https://doi.org/10.3114/sim0003>

514 Bogale, M., Orr, M. -J., O'Hara, M. J., Untereiner, W. A., 2010. Systematics of *Catenulifera*
515 (anamorphic *Hyaloscyphaceae*) with an assessment of the phylogenetic position of *Phialophora*
516 *hyaline*. *Fungal Biology*, 114, 369-409. <https://doi.org/10.1016/j.funbio.2010.02.006>

517 Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C., Al-Ghalith, G. A., Alexander,
518 H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A.,
519 Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, J., Cope, E.
520 K., Da Silva, R., Diener, C., Dorrestein, P. C., Douglas, G. M., Durall, D. M., Duvallet, C.,
521 Edwardson, C. F., Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J. M., Gibbons, S. M., Gibson,
522 D. L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H., Huttenhower, C.,
523 Huttley, G. A., Janssen, S., Jarmusch, A. K., Jiang, L., Kaehler, B. D., Kang, K. B., Keefe, C. R.,
524 Keim, P., Kelley, S. T., Knights, D., Koester, I., Kosciulek, T., Kreps, J., Langille, M. G. I., Lee,
525 J., Ley, R., Liu, Y. -X., Loftfield, E., Lozupone, C., Maher, M., Marotz, C., Martin, B. D.,
526 McDonald, D., McIver, L. J., Melnik, A. V., Metcalf, J. L., Morgan, S. C., Morton, J. T., Naimey,
527 A. T., Navas-Molina, J. A., Nothias, L. F., Orchanian, S. B., Pearson, T., Peoples, S. L., Petras,
528 D., Preuss, M. L., Pruesse, E., Rasmussen, L. B., Rivers, A., Robeson, M. S., Rosenthal, P.,
529 Segata, N., Shaffer, M., Shiffer, A., Sinha, R., Song, S. J., Spear, J. R., Swafford, A. D.,
530 Thompson, L. R., Torres, P. J., Trinh, P., Tripathi, A., Turnbaugh, P. J., Ul-Hasan, S., van der
531 Hooft, J. J. J., Vargas, F., Vázquez-Baeza, Y., Vogtmann, E., von Hippel, M., Walters, W., Wan,
532 Y., Wang, M., Warren, J., Weber, K. C., Williamson, C. H. D., Willis, A. D., Xu, Z. Z., Zaneveld,

533 J. R., Zhang, Y., Zhu, Q., Knight, R., & Caporaso, J. G. (2019). Reproducible, interactive,
534 scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology*, 37, 852-
535 857. <https://doi.org/10.1038/s41587-019-0209-9>

536 Bozoudi, D., & Tsaltas, D. (2018). The multiple and versatile roles of *Aureobasidium pullulans* in
537 the vitivinicultural sector. *Fermentation*, 4(4), 85. <https://doi.org/10.3390/fermentation4040085>

538 Bremner, J. M. (1996). Nitrogen total. In D. Sparks, A. Page, P. Helmke, R. Loeppert, P. N.
539 Soltanpour, M. A. Tabatabai, C. T. Johnston, & M. E. Sumner (Eds.), *Methods of soil analysis*,
540 *Part 3. Chemical Methods SSSA Book Series No 5* (pp. 1085-1121). SSSA and ASA, Madison,
541 WI. <https://doi.org/10.2136/sssabookser5.3.c37>

542 Brindley, G.W., & Brown, G. (1980). Crystal structures of clay minerals and their X-ray
543 identification. *Mineralogic Society Monograph No. 5*. Mineral. Soc. of Great Britain and Ireland,
544 London. <https://doi.org/10.1180/mono-5>

545 Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016).
546 DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, 13,
547 581–583. <https://doi.org/10.1038/nmeth.3869>

548 Chamnipa, N., Thanonkeo, S., Klanrit, P., & Thanonkeo, P. (2018). The potential of the newly
549 isolated thermotolerant yeast *Pichia kudriavzevii* RZ8-1 for high-temperature ethanol production.
550 *Brazilian Journal of Microbiology*, 49, 378-391. <https://doi.org/10.1016/j.bjm.2017.09.002>

551 Chaúque, F. R., Cordani, U. G., & Jamal, D. L. (2019). Geochronological systematics for the
552 Chimoio-Macossa frontal nappe in central Mozambique: Implications for the tectonic evolution of
553 the southern part of the Mozambique belt. *Journal of African Earth Sciences*, 150, 47-67.
554 <https://doi.org/10.1016/j.jafrearsci.2018.10.013>

555 Cheshire, M. V., Christensen, B. T., & Sorensen, L. H. (1990). Labelled and native sugars in particle-
556 size fractions from soils incubated with ¹⁴C straw for 6 to 18 years. *European Journal of Soil*
557 *Science*, 41, 29-39. <https://doi.org/10.1111/j.1365-2389.1990.tb00042.x>

- 558 Chen, L., Xiang, W., Wu, H., Ouyang, S., Zhou, B., Zeng, Y., Chen, Y., & Kuzyakov, Y. (2019).
559 Tree species identity surpasses richness in affecting soil microbial richness and community
560 composition in subtropical forests. *Soil Biology and Biochemistry*, *130*, 113-121.
561 <https://doi.org/10.1016/j.soilbio.2018.12.008>
- 562 Chovanova, K., & Zámocký, M. (2016). Detection of the antibacterial effect of *Chaetomium*
563 *cochliodes* Palliser CCM F-232 based on agar plugs and unprocessed fungal substances from
564 cultivation media. *Biologia*, *71*, 1204–1211. <https://doi:10.1515/biolog-2016-0153>
- 565 Chowdhury, F. I., Barua, I., Chowdhury, A. I., Resco de Dios, V., & Alam, M. S. (2020). Agroforestry
566 shows higher potential than reforestation for soil restoration after slash-and-burn: a case study
567 from Bangladesh. *Geology, Ecology, and Landscape*, *6(1)*, 48-54.
568 <https://doi.org/10.1080/24749508.2020.1743528>
- 569 Climate-data (2019). Retrieved from <https://en.climate-data.org/>.
- 570 Clocchiatti, A., Hannula, S. E., van den Berg, M., & Korthals, G. (2020). The hidden potential of
571 saprotrophic fungi in arable soil: Patterns of short-term stimulation by organic amendments.
572 *Applied Soil Ecology*, *147*, 103434. <https://doi.org/10.1016/j.apsoil.2019.103434>
- 573 Curtis, P. G., Slay, C. M., Harris, N. L., Tyukavina, A., & Hansen, M. C. (2018). Classifying drivers
574 of global forest loss. *Science*, *361*, 1108-1111. <https://doi.org/10.1126/science.aau3445>
- 575 Das, S., Singh, S., Verma, J. P., & Mukherjee, A. (2021). Chapter 14 - Fungi: A potential candidate
576 for sustainable agriculture and agroecosystem. In V. Jay Prakash, C. Macdonald, V. Gupta, & A.
577 Appa Podile (Eds.), *New and Future Developments in Microbial Biotechnology and*
578 *Bioengineering* (pp. 159-164). Elsevier. <https://doi.org/810.1016/B978-0-444-64325-4.00014-6>
- 579 Deaver, N. R., Hesse, C., Kuske, C. R., & Porras-Alfaro, A. (2019). Presence and distribution of
580 insect-associated and entomopathogenic fungi in a temperate pine forest soil: An integrated
581 approach. *Fungal Biology*, *123*, 864-874. <https://doi.org/10.1016/j.funbio.2019.09.006>

582 Deveautour, C., Donn, S., Power, S. A., Bennett, A. E., & Powell, J. R. (2018). Experimentally altered
583 rainfall regimes and host root traits affect grassland arbuscular mycorrhizal fungal communities.
584 *Molecular Ecology*, 27, 2152-2163. <https://doi.org/10.1111/mec.14536>

585 Diao, Y. Z., Chen, Q., Jiang, X. Z., Houbraken, J., Barbosa, R. N., Cai, L., & Wu, W. P. (2019).
586 *Penicillium* section *Lanata-divaricata* from acidic soil. *Cladistics*, 35, 514-249.
587 <https://doi.org/10.1111/cla.12365>

588 Dixon, J. B., & Schulze, S. G. (Eds.), 2002. Soil Mineralogy with Environmental Applications Num.
589 7. In *Soil Science Society of America Book Series*. Soil Sci. Soc. Am. Inc., Madison, Wisconsin,
590 USA.

591 Efthymiou, A., Grønlund, M., Müller-Stöver, D. S., & Jakobsen, I. (2018a). Augmentation of the
592 phosphorus fertilizer value of biochar by inoculation of wheat with selected *Penicillium* strains.
593 *Soil Biology and Biochemistry*, 116, 139-147. <https://doi.org/10.1016/j.soilbio.2017.10.006>

594 Efthymiou, A., Jensen, B., & Jakobsen, I. (2018b). The roles of mycorrhiza and *Penicillium*
595 inoculants in phosphorus uptake by biochar-amended wheat. *Soil Biology and Biochemistry*, 127,
596 168-177. <https://doi.org/10.1016/j.soilbio.2018.09.027>

597 Frac, M., Hannula, S. E., Belka, M., & Jędrzycka, M. (2018). Fungal biodiversity and their role in
598 soil health. *Frontiers in Microbiology*, 9, 1-19. <https://doi.org/10.3389/fmicb.2018.00707>

599 Gay-des-Combes, J. M., Sanz Carrillo, C., Jozef, B., Robroek, M., Jasey, V. E. J., Mills, R. T. E.,
600 Arif, M. S., Falquet, L., Frossard, E., & Buttler, A. (2017). Tropical soils degraded by slash-and-
601 burn cultivation can be recultivated when amended with ashes and compost. *Ecology and*
602 *Evolution*, 7, 5378-5388. <https://doi.org/10.1002/ece3.3104>

603 Giraldo, A., & Crous, P. W. (2019). Inside *Plectosphaerellaceae*. *Studies in Mycology*, 92, 227-286.
604 <https://doi.org/10.1016/j.simyco.2018.10.005>

605 Gizaw, B., Tsegay, Z., Tefera, G., Aynalem, E., Wassie, M., & Abatneh, E. (2017). Phosphate
606 solubilizing fungi isolated and characterized from Teff rhizosphere soil collected from North

607 Showa zone, Ethiopia. *African Journal of Microbiology Research*, 11, 687-696.
608 <https://doi.org/10.5897/AJMR2017.8525>

609 Gonçalves Lintemani, M., Loss, A., Sepulveda Mendes, C., & Fantini, A. C. (2019). Long fallows
610 allow soil regeneration in slash-and-burn agriculture. *Journal of the Science of Food and*
611 *Agriculture*, 100, 1142-1154. <https://doi.org/10.1002/jsfa.10123>

612 Goncharov, A., Glebova, A. A., & Tiunov, A. V. (2020). Trophic interactions between *Fusarium*
613 species and soil fauna: A meta-analysis of experimental studies. *Applied Soil Ecology*, 145,
614 103302. <https://doi.org/10.1016/j.apsoil.2019.06.005>

615 Graystone, S. J., & Campbell, C. D., 1996. Functional biodiversity of microbial community in the
616 rhizospheres of hybrid larch (*Larix europaeis*) and Sitka spruce (*Picea sitchensis*). *Tree*
617 *Physiology*, 16, 1031-1038. <https://doi.org/10.1093/treephys/16.11-12.1031>

618 Habtewold, J. Z., Helgason, B. L., Yanni, S. F., Janzen, H., Ellert, B. H., & Gregorich, E. G. (2020).
619 Litter composition has stronger influence on the structure of soil fungal than bacterial
620 communities. *European Journal of Soil Biology*, 98, 103190.
621 <https://doi.org/10.1016/j.ejsobi.2020.103190>

622 Hamayun, M., Khan, S. A., Khan, A. L., Tang, D. -S., Hussain, J., Ahmad, B., Anwar, Y., & Lee, I.
623 -J. (2010). Growth promotion of cucumber by pure cultures of gibberellin-producing *Phoma* sp.
624 GAH7. *World Journal of Microbiology and Biotechnology*, 26, 889-894.
625 <https://doi.org/10.1007/s11274-009-0248-3>

626 Hosoya, T. (2002). *Hyaloscyphaceae* in Japan (6)**: the genus *Hyphodiscus* in Japan and its
627 anamorph *Catenulifera* gen. nov. *Mycoscience*, 43, 47-57.
628 <https://doi.org/10.1007/s102670200008>

629 Hui, F., Ren, Y., Chen, L., Li, Y., Zhang, L., & Niu, Q. (2014). Molecular phylogenetic analysis
630 reveals the new genus *Hemisphaericaspora* of the family Debaryomycetaceae. *PLoS ONE*, 9,
631 e103737. <https://doi:10.1371/journal.pone.0103737>

- 632 Ji, S., Liu, Z., Liu, B., Wang, Y., Wang, J. (2020). The effect of *Trichoderma* biofertilizer on the
633 quality of flowering Chinese cabbage and the soil environment. *Scientia Horticulturae*, 262,
634 109069. <https://doi.org/10.1016/j.scienta.2019.109069>
- 635 Jumpponen, A., Jones, K. L., Blair, B. (2010). Vertical distribution of fungal communities in tallgrass
636 prairie soil. *Mycologia*, 102, 1027-1041. <https://doi.org/10.3852/09-316>
- 637 Khan, A. L., Al-Harrasi, A., Al-Rawahi, A., Al-Farsi, Z., Al-Mamari, A., Waqas, M., Asaf, S.,
638 Elyassi, A., Mabood, F., Shin, J. H., & Lee, I. J. (2016). Endophytic fungi from frankincense tree
639 improves host growth and produces extracellular enzymes and indole acetic acid. *PLoS ONE*, 11,
640 1-19. <https://doi.org/10.1371/journal.pone.0158207>
- 641 Khan, S. A., Hamayun, M., Khan, A. L., Lee, I. -J., Shinawari, Z. K., & Kom, J. -G. (2012). Isolation
642 of plant growth promoting endophytic fungi from dicots inhabiting coastal sand dunes of Korea.
643 *Pakistan Journal Botany*, 44, 1453-1460.
- 644 Kim, J. S., Baek, J. H., Park, N. -H., & Kim, C. (2016). Complete genome sequence of halophilic
645 yeast *Meyerozyma caribbica* MG20W isolated from rhizosphere soil. *Genome Announcement*, 3,
646 e00127-15. <https://doi:10.1128/genomeA.00127-15>
- 647 Kumla, J., Nundaeng, S., Suwannarach, N., & Lumyong, S. (2020). Evaluation of multifarious plant
648 growth promoting trials of yeast isolated from the soil of assam tea (*Camellia sinensis* var.
649 assamica) plantations in Northern Thailand. *Microorganisms*, 8, 1-18.
650 <https://doi.org/10.3390/microorganisms8081168>
- 651 Lavkulich, L. M., & Wiens, J. H. (1970). Comparison of organic matter destruction by hydrogen
652 peroxide and sodium hypochlorite and its effect on selected mineral constituents. *Soil Science*
653 *Society of America Journal*, 34, 755-758.
654 <https://doi.org/10.2136/sssaj1970.03615995003400050025x>
- 655 Leguina, A. C. del V., Barrios, A. C., del M. Rosales Soro, M., Lacosegliaz, M. J., Pajot, H. F.,
656 Castellanos de Figueroa, L. I., & Nieto-Peñalver, C. G. (2019). Copper alters the physiology of

657 tomato rhizospheric isolates of *Papiliotrema laurentii*. *Scientia Horticulturae*, 243, 376-384.
658 <https://doi.org/10.1016/j.scienta.2018.08.057>

659 Lemmens, M., Buerstmayr, H., Krska, R., Schuhmacher, R., Grausgruber, H., & Ruckenbauer, P.
660 (2004). The effect of inoculation treatment and long-term application of moisture on *Fusarium*
661 head blight symptoms and deoxynivalenol contamination in wheat grains. *European Journal of*
662 *Plant Pathology*, 110, 299-308. <https://doi.org/10.1023/B:EJPP.0000019801.89902.2a>

663 Li, X., Wang, J., Zhang, S., Wang, H., Li, X., Li, X., & Zhang, H. (2018). Distribution of fungal
664 endophytes in roots of *Stipa krylovii* across six vegetation types in grassland of northern China.
665 *Fungal Ecology*, 31, 47-53. <https://doi.org/10.1016/j.funeco.2017.11.001>

666 Limtong, S., Srisuk, N., Yongmanitchai, W., Yurimoto, H., & Nakase, T. (2008). *Ogataea*
667 *chonburiensis* sp. nov. and *Ogataea nakhonphanomensis* sp. nov., thermotolerant,
668 methylotrophic yeast species isolated in Thailand, and transfer of *Pichia siamensis* and *Pichia*
669 *thermomethanolica* to the genus *Ogataea*. *Int. J. Syst. Evol. Microb.* 58, 302-307.
670 <https://doi.org/10.1099/ijs.0.65380-0>

671 Lueders, T., Kindler, R., Miltner, A., Friedrich, M. W., & Kaestner, M. (2006). Identification of
672 bacterial micropredators distinctively active in a soil microbial food web. *Applied and*
673 *Environmental Microbiology*, 72, 5342-5348. <https://doi.org/10.1128/AEM.00400-06>

674 Machinet, G. E., Bertrand, I., Chabbert, B., & Recous, S. (2009). Decomposition in soil and chemical
675 change of maize roots with genetic variations affecting cell wall quality. *European Journal of*
676 *Soil Science*, 60, 176-185. <https://doi.org/10.1111/j.1365-2389.2008.01109.x>

677 Makhuvele, R., Ncube, I., Jansen van Rensburg, E. L., & La Grange, D. C. (2017). Isolation of fungi
678 from dung of wild herbivores for application in bioethanol production. *Brazilian Journal of*
679 *Microbiology*, 48, 648-655. <https://doi.org/10.1016/j.bjm.2016.11.013>

680 Maria, R. M., & Yost, R. (2006). A survey of soil fertility status of four agroecological zones of
681 Mozambique. *Soil Science*, 171(11), 902-914.
682 <https://doi.org/10.1097/01.ss.0000228058.38581.de>

683 Meloun, M., Sáníka, M., Němec, P., Křítková, S., & Kupka, K. (2005). The analysis of soil cores
684 polluted with certain metals using the Box-Cox transformation. *Environmental Pollution*, 137,
685 273–280. <https://doi.org/10.1016/j.envpol.2005.01.027>

686 Mertz, O., Leisz, S. J., Heinemann, A., Rerkasem, K., Dressler, T., Dressler, W., Pham, V. C., Vu, K.
687 C., Schmidt-Vogt, D., Colfer, C. J. P., Epprecht, M., Padoch, C., & Potter, L. (2009). Who
688 counts? Demography of swidden cultivators in Southeast Asia. *Human Ecology*, 37, 281–289.
689 <https://doi.org/10.1007/s10745-009-9249-y>

690 Mohammed, A., Kadhim, J. H., & Hasan, A. M. (2019). Laboratory evaluation of entomopathogenic
691 fungi for the control of khapra beetle (*Coleoptera dermestidae*) and their effects on the beetles'
692 fecundity and longevity. *Journal of Agriculture and Urban Entomology*, 35, 1–12.
693 <https://doi:10.3954/1523-5475-35.1.1>

694 Mota-Gutierrez, J., Ferrocino, I., Rantsiou, K., & Cocolin, L. (2019). Metataxonomic comparison
695 between internal transcribed spacer and 26S ribosomal large subunit (LSU) rDNA gene.
696 *International Journal of Food Microbiology*, 290, 132-140.
697 <https://doi.org/10.1016/j.ijfoodmicro.2018.10.010>

698 Mueller, R. C., Balasch, M., & Kuske, C. R. (2014). Contrasting soil fungal community responses to
699 experimental nitrogen addition using the large subunit rRNA taxonomic marker and
700 cellobiohydrolase I functional marker. *Molecular Ecology*, 23, 4406-4417.
701 <https://doi.org/10.1111/mec.12858>

702 Nakayan, P., Shen, F. -T., Hung, M. -H., & Youn, C. -C. (2009). Effectiveness of *Pichia* sp. CC1 in
703 decreasing chemical fertilization requirements of garden lettuce in pot experiments. *Asian*
704 *Journal of Food and Agro-Industry*, Special Issue, S66-S68.

705 Nath, T. K., Jashimuddin, M., Hasan, Md. K., Shahjahan, Md., & Pretty, J. (2016). The sustainable
706 intensification of agroforestry in shifting cultivation areas of Bangladesh. *Agroforestry Systems*,
707 90, 405-416. <https://doi.org/10.1007/s10457-015-9863-1>

708 Nelson, D. W., & Sommers, L. E. (1996). Total Carbon, Organic Carbon, and Organic Matter. In D.
709 L. Sparks, A. Page, P. Helmke, R. Loeppert, P. N. Soltanpour, M. A. Tabatabai, C. T. Johnston,
710 & M. E. Sumner (Eds.), *Methods of Soil Analysis. Part 3. Chemical Methods, SSSA Book Series*
711 *No. 5* (pp. 961-1010). SSSA and ASA, Madison, WI.

712 Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., Schilling, J. S., &
713 Kennedy, P. G. (2016). FUNGuild: An open annotation tool for parsing fungal community
714 datasets by ecological guild. *Fungal Ecology*, *20*, 241-248.
715 <https://doi.org/10.1016/j.funeco.2015.06.006>

716 Oehl, F., Laczko, E., Oberholzer, H. R., Jansa, J., & Egli, S. (2017). Diversity and biogeography of
717 arbuscular mycorrhizal fungi in agricultural soils. *Biological Fertility of Soils*, *53*, 777-797.
718 <https://doi.org/10.1007/s00374-017-1217-x>

719 Olsen, S. R., Cole, C. V., Watanabe, F. S., & Dean, L. A. (1954). Estimation of available phosphorus
720 in soils by extraction with sodium bicarbonate. *USDA circular 939*. U.S. Gov. Print. Office,
721 Washington, D.C.

722 Oskiera, M., Szczech, M., Stębowska, A., Smolińska, U., Bartoszewski, G. (2017). Monitoring of
723 *Trichoderma* species in agricultural soil in response to application of biopreparations. *Biological*
724 *Control*, *113*, 65-72. <https://doi.org/10.1016/j.biocontrol.2017.07.005>

725 Paula, F. S., Tatti, E., Thorn, C., Abram, F., Wilson, J., O'Flaherty, V. (2020). Soil prokaryotic
726 community resilience, fungal colonisation and increased cross-domain co-occurrence in response
727 to a plant-growth enhancing organic amendment. *Soil Biology and Biochemistry*, *149*, 107937.
728 <https://doi.org/10.1016/j.soilbio.2020.107937>

729 Pongcharoen, P., Chawneua, J., & Tawong, W. (2018). High temperature alcoholic fermentation by
730 new thermotolerant yeast strains *Pichia kudriavzevii* isolated from sugarcane field soil.
731 *Agricultural and Natural Research*, *52*, 511-518. <https://doi.org/10.1016/j.anres.2018.11.017>

732 Procópio, L., & Barreto, C. (2021). The soil microbiomes of the Brazilian Cerrado. *Journal of Soils*
733 *and Sediments*, *21*, 2327–2342. <https://doi.org/10.1007/s11368-021-02936-9>

- 734 Rafael, R. B. A., Fernández-Marcos, M. L., Cocco, S., Ruello, M. L., Weindorf, D. C., Cardelli, V.,
735 & Corti, G. (2018). Assessment of potential nutrient release from phosphate rock and dolostone
736 for application in acid soils. *Pedosphere*, 28, 44-58. [https://doi:10.1016/S1002-0160\(17\)60437-](https://doi:10.1016/S1002-0160(17)60437-5)
737 [5](https://doi:10.1016/S1002-0160(17)60437-5)
- 738 Randriamalala, J. R., Randriarimalala, J., Hervé, D., & Carrière, S. M. (2019). Slow recovery of
739 endangered xerophytic thickets vegetation after slash-and-burn cultivation in Madagascar.
740 *Biological Conservation*, 233, 260-267. <https://doi.org/10.1016/j.biocon.2019.03.006>
- 741 Rožek, K., Rola, K., Błaszowski, J., Leski, T., & Zubek, S. (2020). How do monocultures of fourteen
742 forest tree species affect arbuscular mycorrhizal fungi abundance and species richness and
743 composition in soil?. *Forest Ecology and Management*, 465, 18091.
744 <https://doi.org/10.1016/j.foreco.2020.118091>
- 745 Saliou Sarr, P., Sugiyama, A., Boyogueno Begoude, A. D., Yazakia, K., Araki, S., & Nawata, E.
746 (2019). Diversity and distribution of Arbuscular Mycorrhizal Fungi in cassava (*Manihot*
747 *esculenta* Crantz) croplands in Cameroon as revealed by Illumina MiSeq. *Rhizosphere*, 10, 100-
748 147. <https://doi.org/10.1016/j.rhisph.2019.100147>
- 749 Sandoval-Denis, M., Guarro, J., Cano-Lira, J. F., Sutton, D. A., Wiederhold, N. P., de Hoog, G. S.,
750 Abbott, S. P., Decock, C., Sigler, L., & Gené, J. (2016). Phylogeny and taxonomic revision of
751 *Microascaceae* with emphasis on synnematosus fungi. *Studies in Mycology*, 83, 193-233.
752 <https://doi.org/10.1016/j.simyco.2016.07.002>
- 753 Sannino, C., Borruso, L., Smiraglia, C., Bani, A., Mezzasoma, A., Brusetti, L., Turchetti, B., &
754 Buzzini, P. (2020). Dynamics of *in situ* growth and taxonomic structure of fungal communities
755 in Alpine supraglacial debris. *Fungal Ecology*, 44, 100891.
756 <https://doi.org/10.1016/j.funeco.2019.100891>
- 757 Saravanakumar, K., Shanmuga Arasu, V., & Kathiresan, K. (2013). Effect of *Trichoderma* on soil
758 phosphate solubilization and growth improvement of *Avicennia marina*. *Aquatic Botany*, 104,
759 101-105. <https://doi.org/10.1016/j.aquabot.2012.09.001>

760 Schoeneberger, P. J., Wysocki, D. A., Benham, E. C., & Soil Survey Staff (2012). Field book for
761 describing and sampling soils Version 3.0. Natural Resources Conservation Service, National
762 Soil Survey Center, Lincoln, NE.

763 Serrani, D., Ferrocino, I., Garofalo, C., Osimani, A., Corvaglia, M. R., Milanović, V., Aquilanti, L.,
764 Cocco, S., Cardelli, V., Alves Rafael, R. B., Franciosi, E., Tuohy, K., Clementi, F., & Corti, G.
765 (2023). Soil bacteria communities under slash and burn in Mozambique as revealed by
766 metataxonomic approach. *Pedosphere*, 33, 508-520.
767 <https://doi.org/10.1016/j.pedsph.2022.06.053>

768 Shah, F., Nicolás, C., Bentzer, J., Ellström, M., Smits, M., Rineau, F., Canbäck, B., Floudas, D.,
769 Carleer, R., Lackner, G., Braesel, J., Hoffmeister, D., Henrissat, B., Ahrén, D., Johansson, T.,
770 Hibbett, D. S., Martin, F., Persson, P., & Tunlid, A. (2016). Ectomycorrhizal fungi decompose
771 soil organic matter using oxidative mechanisms adapted from saprotrophic ancestors. *New*
772 *Phytologist*, 209, 1705-1719.

773 Sharmah, D., & Jha, D. K. (2014). Diversity of arbuscular mycorrhizal fungi in undisturbed forest,
774 slash-and-burn field, and monoculture forest of Indo-Burma megadiverse region. *Brazilian*
775 *Journal of Botany*, 37, 339-351. <https://doi.org/10.1007/s40415-014-0075-0>

776 Shi, J., Zhang, T., Geng, S., Liang, F., & Wang, T. (2021). Effect of accumulated temperature on
777 flavour and microbial diversity of japonica rice during storage. *Journal of Stored Products*
778 *Research*, 92, 101779. <https://doi.org/10.1016/j.jspr.2021.101779>

779 Shi, Q., Liu, Y., Shi, A., Cai, Z., Nian, H., Hartmann, M., & Lian, T. (2020). Rhizosphere soil fungal
780 communities of aluminum-tolerant and -sensitive soybean genotypes respond differently to
781 aluminum stress in an acid soil. *Frontiers in Microbiology*, 11, 1177.
782 <https://doi:10.3389/fmicb.2020.01177>

783 Silva-Forsberg, M. C., & Fearnside, P. M. (1997). Brazilian Amazonian *caboclo* agriculture: effect
784 of fallow period on maize yield. *Forest Ecology and Management*, 97, 283-291.
785 [https://doi.org/10.1016/S0378-1127\(97\)00070-4](https://doi.org/10.1016/S0378-1127(97)00070-4)

786 Siteo, A. (2004). Miombo woodlands and HIV/AIDS interactions Mozambique country report
787 forestry working paper. Food and Agriculture Organization of the United Nations, Rome, Italy.

788 Soil Survey Staff (2014). Keys to Soil Taxonomy, 12th ed. United States Department in Agriculture
789 Natural Resources Conservation Service, Washington, DC.

790 Soil Survey Staff (2015). Illustrated guide to soil taxonomy, version 2. U.S. Department of
791 Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln,
792 Nebraska.

793 Soytong, K., Kanokmedhakul, S., Kukongviriyapa, V., & Isobe, M. (2001). Application of
794 *Chaetomium* species (*Ketomium*®) spectrum biological review article fungicide. *Fungal*
795 *Diversity*, 7, 1-15.

796 Spurgeon, D. J., Keith, A. M., Schmidt, O., Lammertsma, D. R., & Faber, J. H. (2013). Land-use and
797 land-management change: relationships with earthworm and fungi communities and soil
798 structural properties. *BMC Ecology*, 13, 46. <https://doi.org/10.1186/1472-6785-13-46>

799 Sterflinger, K. (2010). Fungi: Their role in deterioration of cultural heritage. *Fungal Biology Reviews*,
800 24, 47-55. <https://doi.org/10.1016/j.fbr.2010.03.003>

801 Sterflinger, K. (1998). Temperature and NaCl- tolerance of rock-inhabiting meristematic fungi.
802 *Antonie van Leeuwenhoek, Journal of Microbiology*, 74, 271–281.
803 <https://doi.org/10.1023/A:1001753131034>

804 Suh, S. O., & Zhou, J. (2010). Methylophilic yeasts near *Ogataea (Hansenula) polymorpha*: A
805 proposal of *Ogataea angusta* comb. nov. and *Candida parapolyomorpha* sp. nov. *FEMS Yeast*
806 *Research*, 10, 631-638. <https://doi.org/10.1111/j.1567-1364.2010.00634.x>

807 Suzuki, S., Noble, A. D., Ruaysoongnern, S., & Chinabut, N. (2007). Improvement in water-holding
808 capacity and structural stability of a sandy soil in Northeast Thailand. *Arid Land Res. Manag.*
809 21(1), 37-49. <https://doi.org/10.1080/15324980601087430>

810 Thomas, G. W. (1996). Soil pH and Soil Acidity. In D. L. Sparks, A. L. Page, P. A. Helmke, R. H.
811 Loeppert, P. N. Soltanpour, M. A. Tabatabai, C. T. Johnston, M. E. Sumner (Eds). *Methods of*

812 *Soil Analysis. Part 3. Chemical Methods* (pp. 475–490). Soil Science Society of America and
813 American Society of Agronomy, Madison, WI.

814 Thomaz, E. L. (2013). Slash-and-burn agriculture: Establishing scenarios of runoff and soil loss for
815 a five-year cycle. *Agriculture, Ecosystems & Environment*, *168*, 1-6.
816 <https://doi.org/10.1016/j.agee.2013.01.008>

817 Toju, H., Kishida, O., Katayama, N., & Takagi, K. (2016). Networks depicting the fine-scale co-
818 occurrences of fungi in soil horizons. *PLOS ONE*, *11*, e0165987.
819 <https://doi.org/10.1371/journal.pone.0165987>

820 van der Wal, A., Geydan, T. D., Kuyper, T. W., & de Boer, W. (2013). A thready affair: linking
821 fungal diversity and community dynamics to terrestrial decomposition processes. *FEMS*
822 *Microbiological Reviews*, *37*, 477–494. <https://doi.org/10.1111/1574-6976.12001>

823 Vega, F.E. (2018). The use of fungal entomopathogens as endophytes in biological control: a review.
824 *Mycologia*, *110*, 4-30. <https://doi.org/10.1080/00275514.2017.1418578>

825 Volkmann, M., Whitehead, K., Rütters, H., Rullkötter, J., & Gorbushina, A. A. (2003). Mycosporine-
826 glutamicol-glucoside: A natural UV-absorbing secondary metabolite of rock-inhabiting
827 microcolonial fungi. *Rapid Communications in Mass Spectrometry*, *17*, 897-902.
828 <https://doi.org/10.1002/rcm.997>

829 Warcup, J. H. (1951). The ecology of soil fungi. *Transactions of the British Mycological Society*, *34*,
830 376-399. [https://doi.org/10.1016/S0007-1536\(51\)80065-2](https://doi.org/10.1016/S0007-1536(51)80065-2)

831 Weber, C. F., Vilgalys, R., & Kuske, C. R. (2013). Changes in fungal community composition in
832 response to elevated atmospheric CO₂ and nitrogen fertilization varies with soil horizon.
833 *Frontiers in Microbiology*, *4*, 1-14. <https://doi.org/10.3389/fmicb.2013.00078>

834 Wijnhoud, J. D. (1997). Solos e outros recursos naturais da Estação Agrária de Sussundenga, vol. 1:
835 Relatório. *Série da Terra e Água do Instituto Nacional De Investigação Agrónomica*,
836 comunicação n°93.

- 837 Wood, S. L. R., Rhemtulla, J. M., & Coomes, O. T. (2016). Intensification of tropical fallow-based
838 agriculture: Trading-off ecosystem services for economic gain in shifting cultivation landscapes?
839 *Agriculture, Ecosystem and Environment*, 215, 47–56.
840 <http://dx.doi.org/10.1016/j.agee.2015.09.005>
- 841 Yadav, M., Singh, A., Balan, V., Pareek, N., & Vivekanand, V. (2019). Biological treatment of
842 lignocellulosic biomass by *Chaetomium globosporum*: Process derivation and improved biogas
843 production. *International Journal of Biological Macromolecules*, 128, 176-183.
844 <https://doi.org/10.1016/j.ijbiomac.2019.01.118>
- 845 Yang, H., Yuan, Y., Zhang, Q., Tang, J., Liu, Y., & Chen, X. (2011). Changes in soil organic carbon,
846 total nitrogen, and abundance of arbuscular mycorrhizal fungi along a large-scale aridity gradient.
847 *Catena*, 87, 70-77. <https://doi.org/10.1016/j.catena.2011.05.009>
- 848 Yehia, R. S., Osman, G. H., Assaggaf, H., Salem, R., & Mohamed, M. S. M. (2020). Isolation of
849 potential antimicrobial metabolites from endophytic fungus *Cladosporium cladosporioides* from
850 endemic plant *Zygophyllum mandavillei*. *South African Journal of Botany*, 134, 296-302.
851 <https://doi.org/10.1016/j.sajb.2020.02.033>
- 852 Zalar, P., Gostinčar, C., de Hoog, G. S., Uršič, V., Sudhadham, M., & Gunde-Cimerman, N. (2008).
853 Redefinition of *Aureobasidium pullulans* and its varieties. *Studies in Mycology*, 61, 21-38.
854 <https://doi.org/10.3114/sim.2008.61.02>
- 855 Zhang, F., Xu, X., Wang, G., Wu, B., & Xiao, Y. (2020a). *Medicago sativa* and soil microbiome
856 responses to *Trichoderma* as a biofertilizer in alkaline-saline soils. *Applied Soil Ecology*, 153,
857 103573. <https://doi.org/10.1016/j.apsoil.2020.103573>
- 858 Zhang, K. H., Shi, C. F., Chai, C. Y., & Hui, F. L. (2020b). *Mrakia panshiensis* sp. nov. A new
859 member of the *Cystofilobasidiales* from soil in China, and description of the teleomorphic-stage
860 of *M. Arctica*. *MycKeys*, 74, 75-90. <https://doi.org/10.3897/mycokeys.74.53433>

861 Zhang, Y., Hao, X., Alexander, T. W., Thomas, B. W., Shi, X., & Lupwayi, N. Z. (2018). Long-term
862 and legacy effects of manure application on soil microbial community composition. *Biology and*
863 *Fertility of Soils*, 54, 269-283. <https://doi.org/10.1007/s00374-017-1257-2>

864 Zhou, J., Jiang, X., Zhou, B., Zhao, B., Ma, M., Guan, D., Li, J., Sanfeng, C., Cao, F., Shen, D., &
865 Jie, Q. (2016). Thirty-four years of nitrogen fertilization decreases fungal diversity and alters
866 fungal community composition in black soil in northeast China. *Soil Biology and Biochemistry*,
867 95, 135-143. <https://doi.org/10.1016/j.soilbio.2015.12.012>

868 Zwartendijk, B. W., van Meerveld, H. J., Ghimire, C. P., Ravelona, M., Lahitiana, J., Bruijnzeel, L.
869 A. (2020). Soil water- and overland flow dynamics in a tropical catchment subject to long-term
870 slash-and-burn agriculture. *Journal of Hydrology*, 582, 124287.
871 <https://doi.org/10.1016/j.jhydrol.2019.124287>

872

873 **Figure captions**

874

875 **Fig. 1.** Average-linkage clustering of soil samples based on fungal ASVs' relative abundance at the
876 highest taxonomical rank for locations, land use, and the A and Bo horizons. Manica province,
877 central Mozambique. The color scale represents the ASVs' abundance, denoted as the Z-score,
878 with brown indicating high abundance and blue indicating low abundance.

879 **Fig. 2.** Boxplots showing the differentially abundant fungal ASVs in the soils from Vanduzi,
880 Sussundenga, and Macate. Manica province, central Mozambique. Boxplots with different
881 letters significantly differ at $FDR \leq 0.05$.

882 **Fig. 3.** Correlation between fungi and soil physicochemical properties in the A and Bo horizons of
883 the soils under charcoal kilns, crop fields, and forests at Vanduzi, Sussundenga, and Macate.
884 Manica province, central Mozambique.

885 **Fig. 4.** Boxplots showing the differentially abundant fungi ASVs in the soils under charcoal kilns,
886 crop fields, and forests. Manica province, central Mozambique. Boxplots with different letters
887 significantly differ at $FDR \leq 0.05$.

888 **Fig. 5.** Boxplots showing the differentially abundant fungi ASVs between the A and Bo horizons.
889 Manica province, central Mozambique. Boxplots with different letters significantly differ at
890 $FDR \leq 0.05$.

891